



Cohomology of flag supervarieties and resolutions of determinantal ideals

Steven V Sam and Andrew Snowden

ABSTRACT

We study the coherent cohomology of generalized flag supervarieties. Our main observation is that these groups are closely related to the free resolutions of (certain generalizations of) determinantal ideals. In the case of super Grassmannians, we completely compute the cohomology of the structure sheaf: it is composed of the singular cohomology of a Grassmannian and the syzygies of a determinantal variety. The majority of the work involves studying the geometry of an analog of the Grothendieck–Springer resolution associated with the super Grassmannian; this takes place in the world of ordinary (non-super) algebraic geometry. Our work gives a conceptual explanation of the result of Pragacz–Weyman that the syzygies of determinantal ideals admit an action of the general linear supergroup. In a subsequent paper, we will treat other flag supervarieties in detail.

1. Introduction

Super geometry is a compelling generalization of algebraic geometry, with important connections to physics and pure mathematics. However, a number of fundamental objects in super geometry are still poorly understood. For example, the cohomology of natural vector bundles on flag supervarieties is not known in general. Understanding the cohomology of line bundles would give a super analog of the celebrated Borel–Weil–Bott theorem which, according to [Pen90], was proposed as a general problem by Manin in 1981. In this paper, we develop a general method for attacking this problem in some new cases and use it to completely compute the cohomology of the structure sheaf on the super Grassmannian.

1.1 General approach

Let G be a complex supergroup, let P be a subsupergroup, and let $X = G/P$ (we refer to [MZ11] for generalities on quotients of supergroups). We outline a general approach to study $H^*(X, \mathcal{O}_X)$.

Let G_{ord} , P_{ord} , and X_{ord} be the underlying subschemes of G , P , and X , and let $\mathcal{I}$ be the ideal sheaf defining X_{ord} inside of X . In our cases of interest, G_{ord} is a complex reductive group, P_{ord} is a parabolic subgroup of G_{ord} , the quotient $X_{\text{ord}} = G_{\text{ord}}/P_{\text{ord}}$ is a flag variety (see [MZ11, Proposition 9.3]), $\mathcal{I}/\mathcal{I}^2$ is a locally free coherent sheaf on X_{ord} , and $\text{gr}(\mathcal{O}_X)$ (formed with respect

Received 20 February 2022, accepted in final form 23 February 2023.

2020 *Mathematics Subject Classification* 13D02, 14M30.

Keywords: determinantal ideals, syzygies, supervarieties.

This journal is © [Foundation Compositio Mathematica](#) 2024. This article is distributed with Open Access under the terms of the [Creative Commons Attribution Non-Commercial License](#), which permits non-commercial reuse, distribution, and reproduction in any medium, provided that the original work is properly cited. For commercial re-use, please contact the [Foundation Compositio Mathematica](#).

S. V Sam was supported by NSF grant DMS-1849173. A. Snowden was supported by NSF grant DMS-1453893.

to the $\mathcal{I}$ -adic filtration) is the exterior algebra on $\mathcal{I}/\mathcal{I}^2$. It follows that we have a spectral sequence

$$E_1^{p,q} = H^{p+q}(X_{\text{ord}}, \bigwedge^p(\mathcal{I}/\mathcal{I}^2)) \implies H^{p+q}(X, \mathcal{O}_X).$$

This is our primary tool for connecting the cohomology of supervarieties to ordinary varieties.

To make use of this spectral sequence, we need to understand the cohomology of $\bigwedge(\mathcal{I}/\mathcal{I}^2)$. The genesis of this paper was the observation that the cohomology of exterior algebras, especially on flag varieties, appears in another context: namely, the calculation of syzygies of determinantal varieties (and similar varieties) via the geometric method developed by Kempf, Lascoux, and Weyman, among others (see [Wey03] for an exposition). This allows us to relate the cohomology of $\bigwedge(\mathcal{I}/\mathcal{I}^2)$ to syzygies, in certain cases.

We now explain how this works. Let $\mathfrak{g}$ and $\mathfrak{p}$ be the Lie superalgebras of G and P . For a point $x = gP_{\text{ord}}$ of X_{ord} , let $\mathfrak{p}^x$ be the Lie superalgebra of gPg^{-1} . Let Y be the vector bundle over X_{ord} whose fiber over x is $\mathfrak{p}_1^x$, the odd part of $\mathfrak{p}^x$; this is a closed subvariety of $X_{\text{ord}} \times \mathfrak{g}_1$. Let Z be the image of Y in $\mathfrak{g}_1$; explicitly, Z is the union of all G_{ord} -conjugates of $\mathfrak{p}_1$. In many cases, Z is a determinantal variety, or something of a similar flavor. Let $\tilde{Z}$ be the affinization of Y , which is a finite cover of Z . We refer to this ensemble of varieties as the *Grothendieck–Springer theory* for X since it is analogous to the classical Grothendieck–Springer resolution (and contains some instances of it). We emphasize that X_{ord} , Y , Z , and $\tilde{Z}$ are ordinary (not super) varieties.

Combining the above spectral sequence and the geometric method, we obtain the following theorem. It establishes a link between the cohomology of flag supervarieties and syzygies of determinantal-like varieties.

THEOREM 1.1. *Suppose $H^i(Y, \mathcal{O}_Y) = 0$ for $i > 0$. Letting $S = \text{Sym}(\mathfrak{g}_1^*)$, we have a canonical isomorphism*

$$H^q(X_{\text{ord}}, \bigwedge^{p+q}(\mathcal{I}/\mathcal{I}^2)) = \text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_{p+q}$$

and a spectral sequence

$$E_1^{p,q} = \text{Tor}_{-q}^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_p \implies H^{p+q}(X, \mathcal{O}_X).$$

1.2 The case of Grassmannians

Now suppose that $X = \mathbf{Gr}_{r|s}(\mathbf{C}^{n|m})$ is the super Grassmannian. We apply the method discussed above to study $H^*(X, \mathcal{O}_X)$. Since $\mathbf{Gr}_{r|s}(\mathbf{C}^{n|m}) \cong \mathbf{Gr}_{s|r}(\mathbf{C}^{m|n})$, we may assume, without loss of generality, that $r \geq s$. We summarize some of the key points here.

Let $V_0 = \mathbf{C}^n$ and $V_1 = \mathbf{C}^m$, and put

$$W_0 = \text{Hom}(V_0, V_1), \quad W_1 = \text{Hom}(V_1, V_0), \quad W = W_0 \times W_1.$$

Thus W is identified with the odd part of the Lie superalgebra $\mathfrak{gl}_{n|m}$. The space Y defined in § 1.1 can be described as follows: a point corresponds to a tuple (f, g, R_0, R_1) , where

- R_0 is an r -dimensional subspace of V_0 ,
- R_1 is an s -dimensional subspace of V_1 ,
- $f: V_0 \rightarrow V_1$ is a linear map satisfying $f(R_0) \subset R_1$,
- $g: V_1 \rightarrow V_0$ is a linear map satisfying $g(R_1) \subset R_0$.

The projection map $Y \rightarrow W$ takes (f, g, R_0, R_1) to (f, g) . The only condition on points in the image is that the nullity of f must be at least $r - s$. Hence $Z = Z_0 \times W_1$, where Z_0 is the determinantal variety consisting of linear maps with nullity at least $r - s$; if $n - m \geq r - s$, then $Z_0 = W_0$ and hence $Z = W$.

It turns out that the affinization $\tilde{Z}$ of Y is typically not Z ; that is, there are global functions on Y that do not factor through Z . To see this, suppose that (f, g, R_0, R_1) is a point of Y . Then fg is an operator on V_1 that preserves the subspace R_1 . Thus the characteristic polynomial of $fg|_{R_1}$ is a factor of the characteristic polynomial of fg , and its coefficients give additional global functions on Y . We show that these generate all the additional functions on Y ; this is a non-trivial theorem. From this, we see that $\tilde{Z}$ can be described as the space of tuples (f, g, p) where $f \in Z_0$, $g \in Z_1$, and p is an appropriate factor of the characteristic polynomial of fg .

We prove a number of results about the geometry of this situation. Notably, we show that $\tilde{Z}$ has rational singularities, which allows us to show that $H^i(Y, \mathcal{O}_Y) = 0$ for $i > 0$. We also show that $\mathcal{O}_{\tilde{Z}}$ is a free $\mathcal{O}_Z$ -module of finite rank and that $\mathcal{O}_{\tilde{Z}} \otimes_{\mathcal{O}_Z} \mathbf{C}$ is the singular cohomology ring of a certain Grassmannian. Thus the free resolution of $\mathcal{O}_{\tilde{Z}}$ over $S = \text{Sym}(W^*)$ can be determined from that of $\mathcal{O}_Z$, which was explicitly computed by Lascoux [Las78]. Applying Theorem 1.1, we obtain a spectral sequence computing $H^*(X, \mathcal{O}_X)$, where the terms are composed of the resolution of $\mathcal{O}_Z$ and the singular cohomology of a Grassmannian. Using Lascoux's work, we show that this spectral sequence degenerates. The final result is the following theorem.

THEOREM 1.2. *Let $X = \mathbf{Gr}_{r|s}(\mathbf{C}^{n|m})$ with $r \geq s$. We have the following:*

- (a) Suppose $n - m \geq r - s$.
 - (i) The algebra $H^*(X, \mathcal{O}_X)$ is naturally isomorphic to $H_{\text{sing}}^*(\mathbf{Gr}_s(\mathbf{C}^m), \mathbf{C})$ as a graded algebra.
 - (ii) The $\mathbf{GL}_{n|m}$ -action on $H^*(X, \mathcal{O}_X)$ is trivial.
- (b) Suppose $r - s \geq n - m$, and let $A^* = H_{\text{sing}}^*(\mathbf{Gr}_s(\mathbf{C}^{n+s-r}), \mathbf{C})$.
 - (i) We have a natural isomorphism $H^*(X, \mathcal{O}_X)^{\mathbf{GL}_{n|m}} = A^*$ of graded algebras.
 - (ii) There is a graded $\mathbf{GL}_{n|m}$ -representation E^* such that $H^*(X, \mathcal{O}_X)$ is isomorphic to $A^* \otimes E^*$ as a graded $\mathbf{GL}_{n|m}$ -equivariant A^* -module.
 - (iii) Let $Z_0 \subset \text{Hom}(\mathbf{C}^n, \mathbf{C}^m)$ be the determinantal variety consisting of linear maps of rank at most $n + s - r$, and regard $\mathcal{O}_{Z_0}$ as a quotient of $S = \text{Sym}(\mathbf{C}^n \otimes (\mathbf{C}^m)^*)$. Then we have an isomorphism of $(\mathbf{GL}_n \times \mathbf{GL}_m)$ -representations

$$E^i = \bigoplus_{p \geq 0} \text{Tor}_p^S(\mathcal{O}_{Z_0}, \mathbf{C})_{i+p},$$

where the subscript on Tor denotes the appropriate graded piece. That is, E^i is the i th linear strand of the free resolution of $\mathcal{O}_{Z_0}$.

We note that in case (b) above, $Z_0 = \text{Hom}(\mathbf{C}^n, \mathbf{C}^m)$ if $r = s$ (and $m \geq n$). In particular, $E^i = 0$ for $i \neq 0$ and $E^0 = \mathbf{C}$, and so this situation behaves like case (a).

1.3 Supergroup representations on syzygies

In 1985, Pragacz and Weyman [PW85] discovered that the free resolutions of determinantal ideals carry a natural (but very much non-obvious) representation of a general linear Lie superalgebra. Their construction was generalized and simplified in some subsequent work of Akin–Weyman [AW97, AW00, AW07] and Raicu–Weyman [RW19]. Theorem 1.2 leads to a much more direct and conceptual construction of this action. Indeed, by considering the case where $s = 0$, we find that the i th linear strand of the resolution of a determinantal variety is identified with $H^i(X, \mathcal{O}_X)$ for an appropriate super Grassmannian X , and this obviously carries an action of the super general linear group (as it acts on X). See § 6.5 for more details.

The first author generalized the results of Pragacz–Weyman to (skew-)symmetric matrices in [Sam14]. Here the resolution carries an action of the periplectic superalgebra. The methods of this paper also apply to that case. The details will appear in an upcoming paper by the authors.

Rather than compute syzygies of determinantal ideals with respect to a polynomial ring, one can consider them with respect to an intermediate quotient ring. Using a certain class of complete intersections as such an intermediate quotient ring, a third class of superalgebra actions was discovered in [Sam16] using the orthosymplectic Lie superalgebra. These representations are infinite-dimensional and do not appear to fall into the mold of the current paper. However, one can attempt to treat the orthosymplectic Lie superalgebra using the methods of this paper; see the comments in the next subsection.

1.4 Further work

We have used Theorem 1.1 to compute the cohomology of various other flag supervarieties. The details of this work will appear in a subsequent paper. We summarize some of the results here:

- Let X be the partial flag supervariety parametrizing flags $V_1 \subset \cdots \subset V_k$ in $\mathbf{C}^{n|m}$ where $\dim(V_i) = r_i|s_i$. Suppose $n \geq m$ and $n - m \geq r_i - s_i$ for each i . Then we compute $H^*(X, \mathcal{O}_X)$ completely. The result is similar to Theorem 1.2(a) in that no interesting syzygies appear.
- Let V be a super vector space of dimension $n|n$ equipped with a periplectic form, and let X be the Grassmannian parametrizing totally isotropic subspaces of V of dimension $r|n - r$. Then we compute $H^*(X, \mathcal{O}_X)$ completely, and the result is analogous to Theorem 1.2. In this case, the cohomology is composed of the singular cohomology of an isotropic Grassmannian for a symplectic group and the syzygies of a determinantal ideal of (skew-)symmetric matrices. This result gives a conceptual explanation of the work of the first author [Sam14] that these syzygies carry a representation of the periplectic supergroup.
- Consider an isotropic super Grassmannian X associated with the orthosymplectic supergroup. In this case, the variety Z is a somewhat mysterious analog of a determinantal variety that we have not previously encountered. Our results in this case are more limited, as we do not know the syzygies of Z in general, and there are examples where the spectral sequence computing $H^*(X, \mathcal{O}_X)$ does not degenerate.
- Let X be a Grassmannian associated with the isomeric (also known as type Q, see [NSS22, § 1.5]) supergroup. In this case, X_{ord} is an ordinary Grassmannian, and $\mathcal{I}/\mathcal{I}^2$ is the sheaf of 1-forms on X_{ord} . We thus see that $H^*(X, \bigwedge(\mathcal{I}/\mathcal{I}^2))$ is the de Rham cohomology of X_{ord} and the map $Y \rightarrow Z$ is the usual Grothendieck–Springer map for the general linear Lie algebra. This gives a very direct connection between $H^*(X, \mathcal{O}_X)$ and the singular cohomology of X_{ord} . We can also handle partial flag varieties associated with the isomeric supergroup.
- We can also treat some exceptional supergroups. For example, if X is a complete flag supervariety associated with $D(2, 1; \alpha)$, then we compute $H^*(X, \mathcal{O}_X)$ completely. The variety Z in this case is the zero locus of the hyperdeterminant on $(\mathbf{C}^2)^{\otimes 3}$.

There is one other direction coming out of Theorem 1.2 that we are currently pursuing. Let $X = \mathbf{Gr}_{r|s}(\mathbf{C}^{\infty|\infty})$ be the infinite super Grassmannian (still with $r \geq s$), and let $\mathcal{R}$ and $\mathcal{Q}$ be the tautological subbundle and quotient bundle on X . A natural problem is to study the cohomology of the equivariant bundle $\mathcal{E} = \mathbf{S}_\lambda(\mathcal{Q}) \otimes \mathbf{S}_\mu(\mathcal{R})$, where $\mathbf{S}_\lambda$ denotes a Schur functor. Theorem 1.2 implies that $H^*(X, \mathcal{O}_X)$ is isomorphic to a polynomial ring $\mathbf{C}[c_1, \dots, c_s]$ in this case, where $r \geq s$ and c_i has degree $2i$. We thus see that $H^*(X, \mathcal{E})$ is a module over this polynomial ring. In an upcoming paper, we study $H^*(X, \mathcal{E})$ from this perspective. The module structure allows us to

prove interesting results about the cohomology groups and offers a kind of explanation for the failure of the classical Borel–Weil–Bott theorem in this setting.

REMARK 1.3. Generalizations of the Borel–Weil–Bott theorem to flag supervarieties appear in the literature: see, for instance, [Cou16, GS10, Pen90, PS97]. However, the existing work (that we are aware of) is of a quite different flavor than this paper and generally does not treat the case of the structure sheaf or the vector bundles considered above, with the exception of super projective space; see [GS10, MS11, Ser96].

1.5 Questions and comments

The results of this paper raise a few issues that deserve further exploration.

- We show that the spectral sequence in Theorem 1.1 degenerates for the super Grassmannian. Our proof is rather special and relies on the fact that the representations appearing in the Tor group of a determinantal ideal are multiplicity-free. Is there a more conceptual reason for this degeneration that does not rely on such explicit combinatorial calculation? One obstacle is that, as noted above, there are examples for the orthosymplectic group in which the spectral sequence does not degenerate.
- Theorem 1.2 shows that the coherent cohomology of a super Grassmannian is related to the singular cohomology of some ordinary Grassmannian. The path between these two objects in our proof is somewhat circuitous. Is there a more direct connection? As noted above, in the case of isomeric Grassmannians, the connection to singular cohomology (through de Rham cohomology) is quite clear.
- Are there other homogeneous bundles on the super Grassmannian whose cohomology can be completely computed using the methods of this paper, such as Ω^1 ? Each bundle still has a filtration and associated graded, but the problem is getting a good description of the cohomology of the resulting sheaf.
- A morphism from a superscheme X to the super Grassmannian is equivalent to the data provided by the pullback of the tautological sequence. Hence, given a subbundle of a trivial superbundle on X , we get a morphism from the coherent cohomology of a super Grassmannian to $H^0(X, \mathcal{O}_X)$. This allows one to define “coherent Chern classes” for superbundles.

1.6 Outline

In § 2, we develop our general method for studying the cohomology of flag supervarieties. The rest of the paper is devoted to carrying out this method in the case of super Grassmannians. In § 3 and § 4, we give some preparatory material needed for this. The heart of the paper is § 5, where we study the Grothendieck–Springer theory associated with the super Grassmannian. Finally, in § 6, we apply this work to compute the cohomology of the super Grassmannian.

1.7 Conventions

Since we treat both superalgebras and ordinary (non-super) algebras, we will reserve the term “algebra” or “ring” to mean an ordinary structure and always use the prefix “super” when it is used. The same applies to other objects, such as groups and schemes.

If X is an affine scheme, we use $\mathcal{O}_X$ to denote both its structure sheaf and the ring of global sections; the meaning will be clear from context.

2. Cohomology of flag supervarieties and syzygies

In this section, we outline a general approach to studying the cohomology of flag supervarieties by connecting them to certain determinantal-like varieties appearing in an analog of Grothendieck–Springer theory.

2.1 Super geometry

We begin by recalling some fundamental notions from super geometry. For our purposes, a (*commutative*) *super algebra* is a $\mathbf{Z}/2$ -graded associative unital $\mathbf{C}$ -algebra R such that $xy = (-1)^{|x||y|}yx$ holds for all homogeneous elements $x, y \in R$, where $|x| \in \mathbf{Z}/2$ denotes the degree of x . We use the term *ordinary algebra* to describe a super algebra concentrated in degree 0. Let R be a super algebra. We let $J = J_R$ be the ideal of R generated by its degree 1 elements. Every element of J is nilpotent, and the quotient R/J is an ordinary algebra. We let $\mathrm{Spec}(R)$ be the topological space $\mathrm{Spec}(R/J)$ equipped with a sheaf of super algebras derived from localizations of R in the usual manner.

A *super scheme* over $\mathbf{C}$ is a topological space equipped with a sheaf of super algebras that is locally isomorphic (in a suitable sense) to a space of the form $\mathrm{Spec}(R)$, where R is a super algebra. We use the term *ordinary scheme* to describe a super scheme for which the structure sheaf is concentrated in degree 0; such an object is a scheme in the classical sense. Let X be a super scheme. We let $\mathcal{J} = \mathcal{J}_X$ be the ideal sheaf generated by the degree 1 elements of $\mathcal{O}_X$. The quotient $\mathcal{O}_X/\mathcal{J}_X$ is a sheaf of ordinary rings, and $(X, \mathcal{O}_X/\mathcal{J}_X)$ is an ordinary scheme. We call this the *underlying ordinary scheme* and denote it by X_{ord} . We let $\mathrm{gr}^n(\mathcal{O}_X) = \mathcal{J}^n/\mathcal{J}^{n+1}$ and $\mathrm{gr}(\mathcal{O}_X) = \bigoplus_{n \geq 0} \mathrm{gr}^n(\mathcal{O}_X)$. Each $\mathrm{gr}^n(\mathcal{O}_X)$ is a quasi-coherent sheaf on X_{ord} , and $\mathrm{gr}(\mathcal{O}_X)$ is a quasi-coherent sheaf of super algebras.

We define a super scheme to be *smooth* if X_{ord} is a smooth algebraic variety, $\mathcal{J}/\mathcal{J}^2$ is a locally free coherent sheaf, and the natural map $\bigwedge(\mathcal{J}/\mathcal{J}^2) \rightarrow \mathrm{gr}(\mathcal{O}_X)$ is an isomorphism. We sometimes use the term “supervariety” when speaking of smooth super schemes. Suppose that X is smooth and irreducible. We define its *dimension* to be the pair $d_0|d_1$, where d_0 is the dimension of X_{ord} and d_1 is the rank of $\mathcal{J}/\mathcal{J}^2$. Given a point $x \in X$, the *cotangent superspace* of x is the super vector space $\mathfrak{m}_x/\mathfrak{m}_x^2$, where $\mathfrak{m}_x$ is the maximal ideal at x ; this is easily seen to have dimension $d_0|d_1$. The *tangent superspace* of x is the dual of the cotangent superspace.

The following proposition, which is immediate from the definitions, is our primary tool for connecting the coherent cohomology of super varieties to ordinary algebraic geometry (here, $H^*(X, \mathcal{O}_X)$ is defined to be the usual sheaf cohomology of $\mathcal{O}_X$ thought of as a sheaf of abelian groups on the underlying topological space of X).

PROPOSITION 2.1. *Let X be a smooth super scheme. There is a natural E_1 spectral sequence*

$$E_1^{p,q} = H^{p+q}(X_{\mathrm{ord}}, \bigwedge^p(\mathcal{J}/\mathcal{J}^2)) \implies H^{p+q}(X, \mathcal{O}_X). \quad (2.1a)$$

2.2 The geometric method

Let X be a complex projective variety, and suppose that

$$0 \rightarrow \xi \rightarrow \epsilon \rightarrow \eta \rightarrow 0$$

is an exact sequence of locally free coherent sheaves on X , with $\epsilon = W^* \otimes \mathcal{O}_X$ globally free. We make the following definitions:

- Let $S = \mathrm{Sym}(W^*)$, a polynomial ring, and $W = \mathrm{Spec}(S)$, an affine space.

- Let $Y = \operatorname{Spec}(\operatorname{Sym}(\eta))$. This is the total space of a vector bundle over X , and a closed subvariety of $X \times W$.
- Let $Z \subset W$ be the image of Y under the projection map $X \times W \rightarrow W$. This is a closed subvariety of W as X is projective.
- Let $\tilde{Z}$ be the affinization of Y ; that is, $\tilde{Z}$ is the affine variety with coordinate ring $\mathcal{O}_{\tilde{Z}} = \Gamma(Y, \mathcal{O}_Y)$. The Stein factorization shows that the map $Y \rightarrow Z$ factors through a finite morphism $\tilde{Z} \rightarrow Z$. In particular, $\mathcal{O}_{\tilde{Z}}$ is a finite S -module.

The following proposition describes how to determine the terms of the minimal free resolution of $\mathcal{O}_{\tilde{Z}}$ in certain situations. We remark that $H^i(Y, \mathcal{O}_Y) = H^i(X, \operatorname{Sym}(\eta))$.

PROPOSITION 2.2. *Suppose $H^i(Y, \mathcal{O}_Y) = 0$ for $i > 0$. Then we have a natural isomorphism of vector spaces*

$$\operatorname{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_{p+q} = H^q(X, \bigwedge^{p+q} \xi)$$

for all $p, q \in \mathbf{Z}$, where the subscript indicates the appropriate graded piece.

Proof. We briefly sketch the proof. Consider the cartesian diagram

$$\begin{array}{ccc} X & \xrightarrow{g'} & X \times W \\ f' \downarrow & & \downarrow f \\ \operatorname{Spec}(\mathbf{C}) & \xrightarrow{g} & W, \end{array}$$

where g' is the zero section and g is the inclusion of 0 in W . Regard $\mathcal{O}_Y$ as a coherent sheaf on $X \times W$. Then we have a natural base change isomorphism [Sta21, Tag 08IB]

$$\operatorname{Lg}^* \operatorname{Rf}_*(\mathcal{O}_Y) = \operatorname{Rf}'_* \operatorname{L}(g')^*(\mathcal{O}_Y). \quad (2.2a)$$

By our assumptions, $\operatorname{Rf}_*(\mathcal{O}_Y) = \mathcal{O}_{\tilde{Z}}$, and so the left side above computes $\operatorname{Tor}_*^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})$. Now, we have a Koszul resolution $\mathcal{O}_W \otimes \bigwedge(\xi) \rightarrow \mathcal{O}_Y$. Since this is a flat resolution, it can be used to compute $\operatorname{L}(g')^*$. We find that $\operatorname{L}(g')^*(\mathcal{O}_Y) = \bigwedge(\xi)$, where $\bigwedge(\xi)$ is the complex with $\bigwedge^k(\xi)$ in cohomological degree $-k$ and all differentials zero. The derived pushforward of this under f' is $\bigoplus_{k \geq 0} H^*(X, \bigwedge^k(\xi))$ (ignoring degrees). We refer to [Wey03, Chapter 5] for additional details. $\square$

REMARK 2.3. The functors $\operatorname{Lg}^* \operatorname{Rf}_*$ and $\operatorname{Rf}'_* \operatorname{L}(g')^*$ are both lax monoidal, and the base change map $\operatorname{Lg}^* \operatorname{Rf}_* \rightarrow \operatorname{Rf}'_* \operatorname{L}(g')^*$ is one of lax monoidal functors. It follows that the two sides of (2.2a) are algebras in the derived category of vector spaces, and the isomorphism is one of algebras. From this, it follows that the isomorphism in Proposition 2.2 is compatible with the multiplicative structures on each side. This compatibility is used at two points in the proof of Theorem 1.2.

2.3 The geometric method in super geometry

Combining Propositions 2.1 and 2.2, we obtain the following important result.

THEOREM 2.4. *Let X be a smooth super variety with X_{ord} projective. Suppose that*

$$0 \rightarrow \mathcal{I}/\mathcal{I}^2 \rightarrow \epsilon \rightarrow \eta \rightarrow 0$$

is an exact sequence of locally free coherent sheaves on X_{ord} , with $\epsilon = W^* \otimes \mathcal{O}_{X_{\operatorname{ord}}}$ globally free. Let $S = \operatorname{Sym}(W^*)$, let $Y = \operatorname{Spec}(\operatorname{Sym}(\eta))$, and let $\tilde{Z}$ be the affinization of Y . Suppose $H^i(Y, \mathcal{O}_Y) = 0$ for $i > 0$. Then we have a natural isomorphism

$$\operatorname{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_{p+q} = H^q(X, \bigwedge^{p+q}(\mathcal{I}/\mathcal{I}^2))$$

and a spectral sequence

$$E_1^{p,q} = \mathrm{Tor}_{-q}^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_p \implies H^{p+q}(X, \mathcal{O}_X).$$

We note one corollary of the theorem. Define the *super dimension* of a super vector space V by $\mathrm{sdim}(V) = \dim(V_0) - \dim(V_1)$, and define the *super Euler characteristic* of a super variety X by

$$\chi(X) = \sum_{i \geq 0} (-1)^i \mathrm{sdim} H^i(X, \mathcal{O}_X).$$

COROLLARY 2.5. *Suppose that we are in the setting of Theorem 2.4. Then $\chi(X)$ is the degree of the map $\tilde{Z} \rightarrow W$ over the generic point of W .*

Proof. From the spectral sequence, we find $\chi(X) = \sum_{p \geq 0} (-1)^p \dim \mathrm{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})$, which is easily seen to coincide with the dimension of the vector space $\mathrm{Frac}(S) \otimes_S \mathcal{O}_{\tilde{Z}}$. $\square$

We note that the corollary implies that $\chi(X) = 0$ if Z is a proper subvariety of W , that is, if $Y \rightarrow W$ is not surjective. In this case, assuming $\mathrm{sdim} H^0(X, \mathcal{O}_X) \neq 0$, we find that $\mathcal{O}_X$ necessarily has higher cohomology. One can prove a number of variants of the above corollary, for example, allowing for cases where $H^i(Y, \mathcal{O}_Y)$ is non-vanishing, or replacing “super dimension” with “super character” in the definition of $\chi(X)$ when there is a group acting.

2.4 Grothendieck–Springer theory for flag supervarieties

Let G be a connected supergroup, let P be a subsupergroup of G , let $X = G/P$, and let $\mathcal{I}$ be the ideal sheaf of X_{ord} in X . We will assume that

- G_{ord} is reductive,
- $P_{\mathrm{ord}} \subseteq G_{\mathrm{ord}}$ is a parabolic subgroup, and
- X is smooth.

We now take a look at what Theorem 2.4 amounts to in this case. We note that this discussion is included only to sketch the general situation and provide context; our main results do not depend on it.

Let $x = gP_{\mathrm{ord}}$ be a complex point of X_{ord} . Let $P^x = {}^gP$ and $P_{\mathrm{ord}}^x = {}^gP_{\mathrm{ord}}$, where ${}^g(-)$ denotes conjugation by g . In this way, X_{ord} can be seen as parametrizing G_{ord} -conjugates of either P or P_{ord} . Let $\mathfrak{g}$, $\mathfrak{p}$, and $\mathfrak{p}^x$ be the Lie superalgebras of G , P , and P^x . The tangent superspace to G/P at x is $\mathfrak{g}/\mathfrak{p}^x$, while the tangent space to $G_{\mathrm{ord}}/P_{\mathrm{ord}}$ at x is $\mathfrak{g}_0/\mathfrak{p}_0^x$. Since $\mathcal{J}/\mathcal{J}^2$ is the conormal bundle to $X_{\mathrm{ord}} \subset X$, we see that the fiber of $\mathcal{J}/\mathcal{J}^2$ at x is $(\mathfrak{g}_1/\mathfrak{p}_1^x)^*$. Let $\epsilon = \mathfrak{g}_1^* \otimes \mathcal{O}_X$, and let η be the locally free coherent sheaf on X_{ord} whose fiber at x is $\mathfrak{p}_1^*$. We then have a short exact sequence

$$0 \rightarrow \mathcal{J}/\mathcal{J}^2 \rightarrow \epsilon \rightarrow \eta \rightarrow 0.$$

So far, we have just explained that there is such a sequence for each fiber. However, since the sheaves are all G_{ord} -equivariant and G_{ord} acts transitively on X_{ord} , it follows that there is such a sequence of the sheaves.

We thus see that we are indeed in the setting of Theorem 2.4. The space W is $\mathfrak{g}_1$. The space Y is the vector bundle over X_{ord} whose fiber at x is $\mathfrak{p}_1^x$. In other words, Y consists of points (x, v) where $x \in X_{\mathrm{ord}}$ and $v \in \mathfrak{p}_1^x$. Recall that $Z \subset \mathfrak{g}_1$ is the image of Y under the projection map $X_{\mathrm{ord}} \times W \rightarrow W$. From the description of Y , we see that $Z = \bigcup_{g \in G_{\mathrm{ord}}} {}^g\mathfrak{p}_1$.

The space $\tilde{Z}$ is more subtle. Let $\mathfrak{h} \subset \mathfrak{p}_0$ be a Cartan subalgebra of $\mathfrak{g}_0$, and let $\mathfrak{W}$ be its Weyl group. We have maps

$$\mathfrak{g}_1 \rightarrow \mathfrak{g}_0 \rightarrow \mathfrak{g}_0 // G_{\text{ord}} \xrightarrow{\cong} \mathfrak{h} // \mathfrak{W},$$

where the first map is self-bracket (that is, $v \mapsto [v, v]$), the second map is the GIT quotient map, and the third is the isomorphism coming from the Chevalley restriction theorem. We note that $\mathfrak{h} // \mathfrak{W}$ is the spectrum of the invariant ring $\text{Sym}(\mathfrak{h}^*)^{\mathfrak{W}}$. As Z is a subvariety of $\mathfrak{g}_1$, we thus obtain a natural map $Z \rightarrow \mathfrak{h} // \mathfrak{W}$. We have similar maps

$$\mathfrak{p}_1^x \rightarrow \mathfrak{p}_0^x \rightarrow \mathfrak{h} // \mathfrak{W}',$$

where $\mathfrak{W}'$ is the subgroup of $\mathfrak{W}$ associated with P_{ord} . Since each element of Y belongs to some $\mathfrak{p}_1^x$, these maps define a map $Y \rightarrow \mathfrak{h} // \mathfrak{W}'$, and this map factors through $\tilde{Z}$ since the target is affine. This leads to a commutative diagram

$$\begin{array}{ccc} \tilde{Z} & \longrightarrow & \mathfrak{h} // \mathfrak{W}' \\ \downarrow & & \downarrow \\ Z & \longrightarrow & \mathfrak{h} // \mathfrak{W}. \end{array} \tag{2.6}$$

As we will see in §5, this diagram is not cartesian in general. However, in all cases we have considered, a slight modification is cartesian.

We thus have the following general plan for studying $H^\bullet(X, \mathcal{O}_X)$:

- (a) Show that $H^i(Y, \mathcal{O}_Y)$ vanishes for $i > 0$.
- (b) Determine the minimal free resolution of $\mathcal{O}_{\tilde{Z}}$ over $S = \text{Sym}(\mathfrak{g}_1^*)$.
- (c) Analyze the spectral sequence in Theorem 2.4.

In the remainder of this paper, we carry out this plan in detail for $X = \mathbf{Gr}_{r|s}(\mathbb{C}^{n|m})$.

3. Splitting rings

In §5, we study Grothendieck–Springer theory in detail for the super Grassmannian. In this section and the next section, we gather some preliminary material that will be needed for that. This section develops the theory of splitting and factorization rings. These have also been considered in [EL05, GSS12, Lak10].

3.1 Splitting rings

Let A be a ring, and let $f = \sum_{i=0}^n a_{n-i} u^i$ be a monic polynomial with coefficients in A (so $a_0 = 1$). We define the *splitting ring* of f , denoted by $\text{Split}_A(f)$, to be the quotient $A[\xi_1, \dots, \xi_n]/I$, where I is the ideal generated by the elements

$$a_i - (-1)^i e_i(\xi_1, \dots, \xi_n), \quad 1 \leq i \leq n,$$

where e_i is the i th elementary symmetric function. Thus $\text{Split}_A(f)$ is the universal quotient of $A[\xi_i]$ in which we have $f(u) = \prod_{i=1}^n (u - \xi_i)$. If A is graded and a_i has degree $2i$, then $\text{Split}_A(f)$ is naturally graded, with ξ_i of degree 2. (We include a factor of 2 here for later convenience.) The symmetric group $\mathfrak{S}_n$ acts on $\text{Split}_A(f)$ by permuting the ξ_i and fixing A . Formation of the splitting ring is compatible with base change: if $A \rightarrow A'$ is a homomorphism and f' is the image

of f under $A[u] \rightarrow A'[u]$, then we have a natural isomorphism

$$\text{Split}_{A'}(f') = A' \otimes_A \text{Split}_A(f).$$

We fix A and f for the remainder of this section and let $B = \text{Split}_A(f)$. We also assume A to be noetherian.

3.2 The universal case

Let $A^{\text{univ}} = \mathbf{Z}[a_1, \dots, a_n]$, let $f^{\text{univ}} \in A^{\text{univ}}[u]$ be the polynomial $u^n + \sum_{i=0}^{n-1} a_{n-i}u^i$, and let B^{univ} be the splitting ring of f^{univ} . The map $\mathbf{Z}[\xi_1, \dots, \xi_n] \rightarrow B^{\text{univ}}$ is surjective and has no kernel since $B^{\text{univ}} \otimes \mathbf{C}$ clearly has Krull dimension n . Thus $B^{\text{univ}} = \mathbf{Z}[\xi_1, \dots, \xi_n]$.

The polynomial $f \in A[u]$ is the image of f^{univ} under a unique ring homomorphism $A^{\text{univ}} \rightarrow A$. Since formation of splitting rings is compatible with base change, we have $B = A \otimes_{A^{\text{univ}}} B^{\text{univ}}$. This can be a useful tool for proving results about general splitting rings.

The above discussion shows that we have a cartesian diagram

$$\begin{array}{ccc} \text{Spec}(B) & \longrightarrow & \text{Spec}(B^{\text{univ}}) \\ \downarrow & & \downarrow \\ \text{Spec}(A) & \longrightarrow & \text{Spec}(A^{\text{univ}}). \end{array}$$

We can identify $\text{Spec}(B^{\text{univ}})$ with $\mathbf{A}^n$ and $\text{Spec}(A^{\text{univ}})$ with the quotient $\mathbf{A}^n // \mathfrak{S}_n$ (which is isomorphic to $\mathbf{A}^n$). Here, $\mathbf{A}^n$ and $\mathfrak{S}_n$ can be identified with the Cartan subalgebra and Weyl group of $\mathfrak{gl}_n$. The above diagram is thus similar to (2.6) (with P_{ord} a Borel subgroup, so that $\mathfrak{W}' = 1$), and this is essentially how splitting rings will be relevant in § 5.

3.3 Basic results

Let $\Delta \in A$ be the discriminant of f . We have $\Delta = \prod_{i \neq j} (\xi_i - \xi_j)$ in B . Recall that a ring homomorphism is *syntomic* if it is flat, of finite presentation, and all of the fibers are local complete intersections; see [Sta21, Tag 00SL].

PROPOSITION 3.1. *We have the following:*

- (a) *As an A -module, B is free of rank $n!$.*
- (b) *The map $A \rightarrow B$ is a syntomic.*
- (c) *If A satisfies Serre's condition (S_k) , then so does B . In particular, if A is Cohen–Macaulay, then so is B .*
- (d) *If Δ is a unit of A , then $A \rightarrow B$ is étale.*
- (e) *If A is reduced and Δ is not a zerodivisor, then B is reduced.*
- (f) *The inclusion $A \rightarrow B$ admits an A -linear splitting.*

Proof. (a) It suffices, by base change, to prove the statement in the universal case where $A = \mathbf{Z}[a_1, \dots, a_n]$ and $f = u^n + \sum_{i=0}^{n-1} a_{n-i}u^i$. This is well known, and there are several proofs, but we include one. The map $A \rightarrow B$ is finite as each ξ_i satisfies the universal polynomial. Since $A \rightarrow B$ is a finite map of polynomial rings of the same dimension, it is flat [Sta21, Tag 00R4]. Therefore, B is projective as an A -module, and thus free, as any projective A -module is free (this statement is easy in this case as everything is graded). The rank can be computed over the generic point, where it is well known to be $n!$.

(b) Suppose that $\mathfrak{p}$ is a prime of A . Then $B \otimes_A \kappa(\mathfrak{p})$ is finite over $\kappa(\mathfrak{p})$ by part (a) and therefore of Krull dimension 0. This ring is a quotient of $\kappa(\mathfrak{p})[\xi_1, \dots, \xi_n]$ by n relations and is therefore a complete intersection. It follows that $A \rightarrow B$ is syntomic.

(c) Since (S_k) is syntomic local [Sta21, Tag 036A], the result follows from part (b).

(d) We have $0 = f(\xi_i)$ and so $0 = f'(\xi_i)d\xi_i$. However, $f'(\xi_i) = \prod_{j \neq i} (\xi_i - \xi_j)$ divides Δ and is therefore a unit. Thus $d\xi_i = 0$. We conclude that $\Omega_{B/A} = 0$. Since B is finite flat over A by part (a), it is therefore étale.

(e) Since A is reduced, it satisfies (R_0) and (S_1) . Thus B satisfies (S_1) by part (c). Since $V(\Delta) \subset \operatorname{Spec}(A)$ has codimension 1 and $A[1/\Delta] \rightarrow B[1/\Delta]$ is étale, it follows that B satisfies (R_0) . Thus B is reduced.

(f) This follows from the existence of the Gysin homomorphism constructed in [Lak10, Theorem 8.1] (it has to be precomposed with multiplication by $\prod_i \xi_i^{n-i}$, which is also A -linear). $\square$

PROPOSITION 3.2. *Suppose that $n!$ is invertible in A . Then B is free of rank 1 as an $A[\mathfrak{S}_n]$ -module.*

Proof. Let $\Lambda = \mathbf{Z}[1/n!]$, and let S^λ be the Specht module for $\mathfrak{S}_n$ over Λ . If M is any $\Lambda[\mathfrak{S}_n]$ -module, then the natural map

$$\bigoplus_{|\lambda|=n} \operatorname{Hom}_{\Lambda[\mathfrak{S}_n]}(S^\lambda, M) \otimes S^\lambda \rightarrow M$$

is an isomorphism. To see this, note that S^λ is a projective $\Lambda[\mathfrak{S}_n]$ -module: Young idempotents are defined over $\Lambda[\mathfrak{S}_n]$ (see [FH04, Lemma 4.26]), so S^λ is a direct summand of $\Lambda[\mathfrak{S}_n]$. Hence both sides above are exact functors of M . It thus suffices to prove the result for $M = S^\mu$ (since these are enough projectives), where it is clear.

Now, it suffices to prove the proposition in the universal case $A = \Lambda[a_1, \dots, a_n]$. By the above isomorphism, we see that $N_\lambda = \operatorname{Hom}_{\Lambda[\mathfrak{S}_n]}(S^\lambda, B)$ is a summand of B and thus projective as an A -module; since N_λ is also graded, it is free. To prove the result, it is enough to show that the A -rank of N_λ coincides with the Λ -rank of S^λ . This can be checked over $\operatorname{Frac}(A)$, where it is well known: $\mathbf{Q}(\xi_1, \dots, \xi_n)$ is isomorphic to the regular representation of $\mathfrak{S}_n$ over $\mathbf{Q}(a_1, \dots, a_n) = \mathbf{Q}(\xi_1, \dots, \xi_n)^{\mathfrak{S}_n}$ by the normal basis theorem in Galois theory. $\square$

REMARK 3.3. The hypothesis that $n!$ is invertible is necessary: indeed, if $n = 2$, $A = \mathbf{F}_2$, and $f = u^2$, then $B \cong \mathbf{F}_2[t]/t^2$ with trivial $\mathfrak{S}_2$ -action, but $\mathbf{F}_2[\mathfrak{S}_2]$ has non-trivial $\mathfrak{S}_2$ -action.

We pause to give a geometric source of splitting rings (see [GSS12, Theorem 6.1]). Let X be a smooth variety over an algebraically closed field, and let $\mathcal{E}$ be a rank n vector bundle on X . Let A be the Chow ring of X , and let $a_i = (-1)^i c_i(\mathcal{E})$, where $c_i(\mathcal{E})$ is the i th Chern class of $\mathcal{E}$. Then the splitting ring B is the Chow ring of the relative flag variety $\mathbf{Fl}(\mathcal{E})$. Furthermore, on $\mathbf{Fl}(\mathcal{E})$, the pullback of $\mathcal{E}$ has a complete flag of subbundles (that is, whose successive quotients are line bundles), and the Chern classes of these line bundles are identified with the $-\xi_i$.

An important case for us is where $X = \operatorname{Spec}(\mathbf{C})$ and $\mathcal{E} = \mathbf{C}^n$, so that $f = u^n$. In that case, this discussion gives the following result (we note that the Chow ring and singular cohomology ring of $\mathbf{Fl}(\mathbf{C}^n)$ are isomorphic since it has a cellular decomposition).

PROPOSITION 3.4. *Suppose $f = u^n$. Regard A as graded and concentrated in degree 0 and B as graded with each ξ_i of degree 2. Then we have a natural isomorphism of graded rings*

$$B = A \otimes H_{\operatorname{sing}}^*(\mathbf{Fl}(\mathbf{C}^n), \mathbf{Z}).$$

3.4 First normality criterion

We now turn our attention to the question of when B is normal. We will give an initial criterion here and a variant in §3.6 that is somewhat more convenient. Let $\tilde{E} \subset \text{Spec}(B)$ be the closed set where at least three of the ξ_i coincide or there are two pairs of ξ_i which coincide, and let $E \subset \text{Spec}(A)$ be the image of $\tilde{E}$. Note that E is closed since $A \rightarrow B$ is finite. We say that an element f of a normal ring R is *squarefree* if $v_{\mathfrak{p}}(f) \in \{0, 1\}$ for all height 1 primes $\mathfrak{p}$ of R , where $v_{\mathfrak{p}}$ denotes the valuation associated with $\mathfrak{p}$. We say that a subset of $\text{Spec}(A)$ has *codimension at least c* if all primes it contains have height at least c .

PROPOSITION 3.5. *Suppose that the following conditions hold:*

- (a) *The ring A is normal.*
- (b) *The discriminant Δ of f is squarefree and not a zerodivisor.*
- (c) *The set E has codimension at least 2 in $\text{Spec}(A)$.*

Then B is normal.

Proof. First suppose that A is a strictly henselian discrete valuation ring. We show that B is regular. If Δ is a unit of A , then B is étale over A and thus regular. Then assume that Δ is not a unit; by condition (b), it is a uniformizer of A . Let $\bar{f}$ be the image of f in $\kappa[u]$, where κ is the residue field of A . Since Δ maps to 0 in κ , it follows that $\bar{f}$ has a repeated root; by condition (c), it has only one (that is, it has $n - 1$ distinct roots). We thus have a factorization $\bar{f} = \bar{q} \cdot \bar{g}$ over κ , where $\bar{q}$ is a degree 2 polynomial with a repeated root over $\bar{\kappa}$ and $\bar{g}$ has distinct roots over $\bar{\kappa}$. Since κ is separably closed, it follows that $\bar{g} = (u - \bar{x}_3) \cdots (u - \bar{x}_n)$ splits completely for $\bar{x}_3, \dots, \bar{x}_n \in \kappa$ distinct (and distinct from the unique root of $\bar{q}$). By the henselian property, we thus have a factorization $f(u) = q(u)(u - x_3) \cdots (u - x_n)$, where $x_i \in A$ lifts $\bar{x}_i$ for $i \geq 3$ and $q(u)$ is a quadratic polynomial with coefficients in A .

Now let $\mathfrak{p}$ be a prime of B above the maximal ideal of A , and work in $B_{\mathfrak{p}}$ in what follows. Applying a permutation if necessary, we can assume $\bar{\xi}_i = \bar{x}_i$ for $i \geq 3$, where $\bar{\xi}_i$ is the image of ξ_i in $B_{\mathfrak{p}}/\mathfrak{p} \cong \kappa$. For $i \geq 3$, it follows that $q(\xi_i)$ and $\xi_i - x_j$, for $j \neq i$, are non-zero in κ , and thus units of $B_{\mathfrak{p}}$; since $f(\xi_i) = 0$, we conclude that $\xi_i = x_i$. This shows that $B_{\mathfrak{p}}$ is generated as an A -algebra by ξ_1 and ξ_2 . We have

$$q(u) \prod_{i \geq 3} (u - \xi_i) = \prod_{i \geq 1} (u - \xi_i).$$

Since monic polynomials are not zerodivisors, it follows that $q(u) = (u - \xi_1)(u - \xi_2)$. From this, we see that $B_{\mathfrak{p}} \cong A[u]/q$.

We consider two cases. If $\bar{q}$ is irreducible (this is only possible if κ has characteristic 2), then $B_{\mathfrak{p}}/\mathfrak{m}B_{\mathfrak{p}}$ is a degree 2 field extension of κ and hence $\mathfrak{m}B_{\mathfrak{p}}$ is the maximal ideal of $B_{\mathfrak{p}}$ which is generated by a single non-nilpotent element, so $B_{\mathfrak{p}}$ is a DVR.

Otherwise, we may assume $\bar{q} = (u - \bar{x}_1)^2$ for some $\bar{x}_1 \in \kappa$. Let π be the discriminant of q . This is an element of A that divides Δ and is 0 in κ , and is therefore a uniformizer. Up to a linear substitution in A , we may assume $\bar{q} = u^2$. Write $q = u^2 + \alpha_1 u + \alpha_2$. Then $\alpha_1, \alpha_2 \in \mathfrak{m}$ but $\alpha_2 \notin \mathfrak{m}^2$ since $\pi = \alpha_1^2 - 4\alpha_2$ is a uniformizer, and hence α_2 is also a uniformizer for A . The maximal ideal of $B_{\mathfrak{p}}$ is generated by u and $\mathfrak{m}B_{\mathfrak{p}}$, but the latter is generated by $\alpha_2 = -(u + \alpha_1)u$, and so just u suffices to generate. Finally, u is not nilpotent since α_2 is not, which shows that $B_{\mathfrak{p}}$ is a DVR.

We now treat the general situation. Let $\mathfrak{p}$ be a height 1 prime of A . We show that $B_{\mathfrak{p}}$ is regular. Let $A_{\mathfrak{p}}^{\text{sh}}$ be the strict henselization of the DVR $A_{\mathfrak{p}}$. By the previous paragraphs, we see

that $B_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} A_{\mathfrak{p}}^{\text{sh}}$ is regular. Now, $A_{\mathfrak{p}}^{\text{sh}}$ is the direct limit of a family $\{A_i\}$ of rings, each of which is an étale cover of $A_{\mathfrak{p}}$. The above arguments apply with A_i in place of $A_{\mathfrak{p}}^{\text{sh}}$ for i sufficiently large. We conclude that $B_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} A_i$ is regular for some i . Since regularity is étale local, we conclude that $B_{\mathfrak{p}}$ is regular. Thus B is regular in codimension 1. Finally, since A is normal, it satisfies Serre's condition (S_2) , and hence the same is true for B by Proposition 3.1(c), so B is therefore normal. $\square$

3.5 Preliminaries on discriminants

In the next subsection, we refine the above normality criterion. Here we prove a few lemmas that will be needed to do this.

LEMMA 3.6. *Let S be a local ring with maximal ideal $\mathfrak{m}$, and let X be an $n \times n$ matrix with entries in S . Suppose that the reduction of X modulo $\mathfrak{m}$ has nullity at least r . Then $\det(X) \in \mathfrak{m}^r$.*

Proof. We argue by induction on n . For $n = 0$ the result is vacuous. Now suppose that the result is known for matrices of size less than n , and let us prove it for matrices of size n .

Suppose that the first column of X has entries in $\mathfrak{m}$. Let X_i be the matrix obtained from X by deleting the first column and the i th row. Then each X_i has nullity at least $r - 1$ modulo $\mathfrak{m}$. Thus, by induction, $\det(X_i) \in \mathfrak{m}^{r-1}$ for each i . Taking the Laplace expansion for $\det(X)$ along the first column, we see that $\det(X) \in \mathfrak{m}^r$.

Now suppose that some entry in the first column of X is a unit. Performing row operations, we can reduce to the case where the first column is the first standard basis vector. Let X' be the matrix obtained by deleting the first row and column of X . Then $\det(X') = \det(X)$. But X' has nullity at least r modulo $\mathfrak{m}$, so $\det(X') \in \mathfrak{m}^r$ by induction. $\square$

We recall some basic facts about Sylvester matrices and discriminants. Let S be a commutative ring, and let $f(u) = a_0 u^n + \cdots + a_n$ and $g(u) = b_0 u^m + \cdots + b_m$ be univariate polynomials with coefficients in S such that $a_0 \neq 0$ (we do not require anything about g and in fact allow the case where it is identically 0). We define their *Sylvester matrix* to be the square matrix of size $n + m$ as follows:

$$\text{Syl}_{n,m}(f, g) = \begin{bmatrix} a_0 & a_1 & a_2 & \cdots & a_n & 0 & 0 & \cdots & 0 \\ 0 & a_0 & a_1 & \cdots & & a_n & 0 & \cdots & 0 \\ \vdots & & & & & & & & \\ 0 & 0 & 0 & & & & & \cdots & a_n \\ b_0 & b_1 & b_2 & \cdots & & & & & \\ \vdots & & & & & & & & \\ 0 & 0 & 0 & & & & & \cdots & b_m \end{bmatrix}.$$

In other words, the first m rows consist of shifts of the sequence $a_0, \dots, a_n$, and the last n rows consist of shifts of the sequence $b_0, \dots, b_m$.

The following is well known, but we include proofs to keep the discussion self-contained.

PROPOSITION 3.7. (a) *If S is a field, then*

$$\deg(\gcd(f, g)) = \dim \ker \text{Syl}_{n,m}(f, g).$$

(b) *If S is graded such that $\deg(a_i) = \deg(b_i) = i$, then $\det(\text{Syl}_{n,m}(f, g))$ is homogeneous of degree mn .*

(c) If $n = \deg(f)$, then the discriminant of f is $\det(\text{Syl}_{n,n-1}(f, f'))$ (up to a sign), where f' is the derivative of f with respect to x .

Proof. (a) Let $\alpha = \alpha_0 u^{m-1} + \cdots + \alpha_{m-1}$ and $\beta = \beta_0 u^{n-1} + \cdots + \beta_{n-1}$. The coefficients of $\alpha f + \beta g$ are the entries of $\begin{bmatrix} \alpha_0 & \cdots & \alpha_{m-1} & \beta_0 & \cdots & \beta_{n-1} \end{bmatrix} \text{Syl}_{n,m}(f, g)$, so that $\ker \text{Syl}_{n,m}(f, g)^T$ is the space of pairs (α, β) with $\deg \alpha < m$ and $\deg \beta < n$ such that $\alpha f + \beta g = 0$. Let $h = \gcd(f, g)$ and $f_0 = f/h$ and $g_0 = g/h$. Then we get $\alpha f_0 = -\beta g_0$, which implies that there is a polynomial γ such that $\alpha = g_0 \gamma$ and $\beta = f_0 \gamma$ and

$$\deg \gamma = \deg \beta - \deg f_0 < \deg h.$$

On the other hand, if γ is any polynomial with $\deg \gamma < \deg h$, then $\alpha = g_0 \gamma$ has degree less than $\deg g \leq m$ and $\beta = f_0 \gamma$ has degree less than $\deg f = n$, and so we have an isomorphism between the polynomials of degree less than $\deg h$ and $\ker \text{Syl}_{n,m}(f, g)$.

(b) Scaling each a_i and b_i by λ^i in $\text{Syl}_{n,m}(f, g)$ is the same as scaling the i th column by λ^{i-1} , the first m rows by $1, \lambda^{-1}, \dots, \lambda^{-(m-1)}$, and the last n rows by $1, \lambda^{-1}, \dots, \lambda^{-(n-1)}$. Hence we see that $\det(\text{Syl}_{n,m}(f, g))$ is homogeneous of degree $\binom{n+m}{2} - \binom{n}{2} - \binom{m}{2} = mn$.

(c) It suffices to prove this in the universal case $S = \mathbf{Z}[a_0, \dots, a_n]$ since both the discriminant of f and $\det(\text{Syl}_{n,n-1}(f, f'))$ are polynomials in the coefficients of f . We grade S by $\deg(a_i) = i$. In this case, both expressions are homogeneous polynomials of degree $n(n-1)$; this holds for $\det(\text{Syl}_{n,n-1}(f, f'))$ by part (b). The discriminant is an irreducible polynomial and vanishes whenever f has a multiple root. By part (a), $\det(\text{Syl}_{n,n-1}(f, f'))$ vanishes whenever $\gcd(f, f')$ has positive degree, which is equivalent to f having a multiple root. Hence the two polynomials agree up to sign. $\square$

LEMMA 3.8. Suppose that S is a local ring with maximal ideal $\mathfrak{m}$, and let $g \in S[u]$ be a monic polynomial of degree n . Suppose that the reduction of g modulo $\mathfrak{m}$ has at most r distinct roots in the algebraic closure of $S/\mathfrak{m}$. Then the discriminant of g belongs to $\mathfrak{m}^{n-r}$.

Proof. Let f be the reduction of g modulo $\mathfrak{m}$. By Lemma 3.6 and Proposition 3.7, the discriminant of g belongs to $\mathfrak{m}^d$, where $d = \deg(\gcd(f, f'))$. For λ in the algebraic closure of $S/\mathfrak{m}$, if $(u - \lambda)^e$ divides f , then $(u - \lambda)^{e-1}$ divides f' , and so $d \geq n - r$. $\square$

3.6 Second normality criterion

We now give a variant of Proposition 3.5. Define $V(\Delta, \partial\Delta) \subset \text{Spec}(A)$ to be the set of points x at which Δ vanishes to order 2, in the sense that it belongs to $\mathfrak{m}_x^2$. If A is finitely generated over a field k and x is a smooth point of $\text{Spec}(A)$, then x belongs to $V(\Delta, \partial\Delta)$ if and only if $\Delta = 0$ in $\kappa(x)$ and $d\Delta = 0$ in $\Omega_{A/k}^1 \otimes_A \kappa(x)$. Since $\Omega_{A/k}^1$ is locally free on the smooth locus, this shows that $V(\Delta, \partial\Delta)$ is closed in the smooth locus. Recall the set $E \subset \text{Spec}(A)$ from § 3.4.

LEMMA 3.9. We have $E \subset V(\Delta, \partial\Delta)$.

Proof. Let $x \in E$, and let $\mathfrak{p} = \mathfrak{p}_x$ be the corresponding prime ideal of A . By the definition of E , the polynomial $f \in A_{\mathfrak{p}}[u]$ has at most $\deg(f) - 2$ distinct roots in the residue field. Thus by Lemma 3.8, we see that $\Delta \in \mathfrak{p}^2 A_{\mathfrak{p}} = \mathfrak{m}_x^2$, and so $x \in V(\Delta, \partial\Delta)$. $\square$

PROPOSITION 3.10. Suppose that the following conditions hold:

- (a) The ring A is normal.
- (b) The discriminant Δ of f is not a zerodivisor.

(c) The set $V(\Delta, \partial\Delta)$ has codimension at least 2.

Then Δ is squarefree and B is normal.

Proof. We apply Proposition 3.5. The set E there has codimension at least 2 by the present assumption (c) and Lemma 3.9. It thus suffices to prove that Δ is squarefree.

Let $\mathfrak{p}$ be a height 1 prime of A . If $V(\mathfrak{p})$ is not an irreducible component of $V(\Delta)$, then $v_{\mathfrak{p}}(\Delta) = 0$. Now suppose that $V(\mathfrak{p})$ is an irreducible component of $V(\Delta)$. Then $v_{\mathfrak{p}}(\Delta) \geq 1$, and we show that equality holds. Suppose toward a contradiction that $v_{\mathfrak{p}}(\Delta) \geq 2$. We can then write $\Delta = ab^k$, where $k \geq 2$, a is a rational function on $\text{Spec}(A)$ that is a unit at $\mathfrak{p}$, and b is a rational function on $\text{Spec}(A)$ that is a uniformizer at $\mathfrak{p}$. Since a is a unit of $A_{\mathfrak{p}}$ and b belongs to $\mathfrak{p}A_{\mathfrak{p}}$, we see that Δ belongs to $\mathfrak{p}^2A_{\mathfrak{p}}$. Thus $\mathfrak{p} \in V(\Delta, \partial\Delta)$, which contradicts assumption (c). We conclude that Δ is squarefree. $\square$

3.7 Factorization rings

Let p and q be non-negative integers such that $p + q = n$, and put $g = \sum_{i=0}^p b_{p-i}u^i$ and $h = \sum_{i=0}^q c_{q-i}u^i$, where $b_0 = c_0 = 1$ and the remaining b_i and c_i are formal symbols. We define the (p, q) -factorization ring of f , denoted by $\text{Fact}_A^{p,q}(f)$, to be $A[b_1, \dots, b_p, c_1, \dots, c_q]/I$, where I is the ideal generated by the elements

$$a_{n-i} - \sum_{j=0}^i b_{p-j}c_{q-i+j}, \quad 0 \leq i \leq n-1.$$

Thus $\text{Fact}_A^{p,q}(f)$ is the universal quotient of $A[b_i, c_j]$ in which we have $f(u) = g(u)h(u)$. It follows from the above formula and the condition $b_0 = 1$ that $\text{Fact}_A^{p,q}(f)$ is generated as an A -algebra by $b_1, \dots, b_p$. If A is graded and a_i is homogeneous of degree $2i$, then $\text{Fact}_A^{p,q}(f)$ is graded and b_i and c_i have degree $2i$. Formation of the factorization ring is compatible with base change, as with the splitting ring.

In what follows, we let $B = \text{Split}_A(f)$ and $C = \text{Fact}_A^{p,q}(f)$.

PROPOSITION 3.11. *We have the following:*

- (a) We have a natural A -algebra isomorphism $B = \text{Split}_C(g) \otimes_C \text{Split}_C(h)$.
- (b) As an A -module, C is free of rank $\binom{n}{p}$.
- (c) The map $A \rightarrow C$ is syntomic.
- (d) If A satisfies Serre's condition (S_k) , then so does C . In particular, if A is Cohen–Macaulay, then so is C .
- (e) If B is reduced (respectively, integral, normal), then C is reduced (respectively, integral, normal).

Proof. (a) Let $\eta_1, \dots, \eta_p$ be the generators of $\text{Split}_C(g)$ and $\eta_{p+1}, \dots, \eta_{p+q}$ those for $\text{Split}_C(h)$. Put $B' = \text{Split}_C(g) \otimes_C \text{Split}_C(h)$. Since $f(u) = \prod_{i=1}^n (u - \eta_i)$ holds over B' , we have an A -algebra homomorphism $\varphi: B \rightarrow B'$ given by $\varphi(\xi_i) = \eta_i$. Let $g^*(u) = \prod_{i=1}^p (u - \xi_i)$ and $h^*(u) = \prod_{i=p+1}^n (u - \xi_i)$ be polynomials in $B[u]$. The factorization $f(u) = g^*(u)h^*(u)$ gives an A -algebra homomorphism $C \rightarrow B$ mapping $g(u)$ to $g^*(u)$ and $h(u)$ to $h^*(u)$. The tautological splittings of $g^*(u)$ and $h^*(u)$ over B yield an A -algebra homomorphism $\psi: B' \rightarrow B$ given by $\psi(\eta_i) = \xi_i$. Since φ and ψ are clearly inverses, the result follows.

(b) By Proposition 3.1(a), we have an A -module isomorphism $B \cong A^{\oplus n!}$ and C -module isomorphisms $\text{Split}_C(g) \cong C^{\oplus p!}$ and $\text{Split}_C(h) \cong C^{\oplus q!}$. Comparing with part (a), we obtain an

A -module isomorphism $C^{\oplus p!q!} \cong A^{\oplus n!}$. It follows that C is projective as an A -module of constant rank $\binom{n}{p}$. As in the proof of Proposition 3.1(a), it follows that C is free in the universal case, and thus in all cases.

(c) Let $\mathfrak{p}$ be a prime of A . Then $C \otimes_A \kappa(\mathfrak{p})$ is finite over $\kappa(\mathfrak{p})$ by part (b), and therefore of Krull dimension 0. This ring is a quotient of $\kappa(\mathfrak{p})[b_1, \dots, b_p, c_1, \dots, c_q]$ by $p + q$ relations and is therefore a complete intersection. Thus $A \rightarrow C$ is syntomic.

(d) This follows since the property is syntomic local.

(e) From part (a), the ring C is isomorphic to a subring of B , which handles the reduced and integral conditions. For the normality condition, we use that B is an iterated splitting ring over C by part (a), and hence the inclusion $C \rightarrow B$ admits a C -linear splitting by Proposition 3.1(f). $\square$

REMARK 3.12. Using Proposition 3.2 and Proposition 3.11(a), we can show that $C = B^{\mathfrak{S}_p \times \mathfrak{S}_q}$ if $n!$ is invertible in A . This gives a C -linear splitting of $C \rightarrow B$ and an alternative proof of Proposition 3.11(e). In fact, the assumption that $n!$ is invertible is relaxed substantially in [EL05, Theorem 3.1]: it suffices to know that either 2 or the discriminant of f is not a zerodivisor in A to show that $C = B^{\mathfrak{S}_p \times \mathfrak{S}_q}$.

As with splitting rings, there is a geometric source of factorization rings (see [GSS12, Theorem 6.1]). Let X be a smooth variety over an algebraically closed field, and let $\mathcal{E}$ be a rank n vector bundle on X . Then the $(p, n - p)$ factorization ring C (again taking A to be the Chow ring of X and $a_i = (-1)^i c_i(\mathcal{E})$) is the Chow ring of the relative Grassmannian $\mathbf{Gr}(p, \mathcal{E})$. When $X = \mathrm{Spec}(\mathbf{C})$, we get the following result.

PROPOSITION 3.13. *Suppose $f = u^n$. Regard A as graded and concentrated in degree 0 and C as graded in the usual manner. Then we have a natural isomorphism of graded rings*

$$C = A \otimes H_{\mathrm{sing}}^*(\mathbf{Gr}_p(\mathbf{C}^n), \mathbf{Z}).$$

REMARK 3.14. One can also form partial splitting rings which are intermediate between B and C , and all of the above properties generalize. For these rings, one sees the cohomology of a partial flag variety in the analog of Proposition 3.13.

4. Some additional preparatory material

In this section, we give a bit more material that will be needed in §5.

4.1 Rational singularities

We first recall some generalities on rational singularities. Let X be an irreducible variety over the complex numbers. We say that X has *rational singularities* if there exists a proper birational map $\pi: Y \rightarrow X$ with Y smooth (that is, a resolution of singularities) such that $\pi_*(\mathcal{O}_Y) = \mathcal{O}_X$ and $R^i\pi_*(\mathcal{O}_Y) = 0$ for $i > 0$. If X has rational singularities, then for any resolution of singularities $\pi': Y' \rightarrow X$, we have $\pi'_*(\mathcal{O}_{Y'}) = \mathcal{O}_X$ and $R^i\pi'_*(\mathcal{O}_{Y'}) = 0$ for $i > 0$; furthermore, X is normal and Cohen–Macaulay [KM98, Theorem 5.10]. We require the following two additional results concerning rational singularities.

PROPOSITION 4.1. *Suppose that X is normal and Cohen–Macaulay, and let $\pi: Y \rightarrow X$ be a resolution of singularities. Let $U \subset X$ be a subset which has rational singularities, and suppose that $\pi^{-1}(X \setminus U)$ has codimension at least 2 in Y . Then X has rational singularities.*

Proof. Since X is Cohen–Macaulay, let ω_X be its dualizing sheaf. We have a natural map $\pi_*\omega_Y \rightarrow \omega_X$ obtained by duality from $\mathcal{O}_X \rightarrow \pi_*\mathcal{O}_Y$, and X has rational singularities if and only if this map is surjective [KM98, Theorem 5.10]. This property is local, so we may assume that X is affine. Consider the following commutative diagram:

$$\begin{array}{ccc} \Gamma(Y, \omega_Y) & \longrightarrow & \Gamma(X, \omega_X) \\ \downarrow & & \downarrow \\ \Gamma(\pi^{-1}(U), \omega_Y) & \longrightarrow & \Gamma(U, \omega_X). \end{array}$$

The bottom map is an isomorphism since U has rational singularities. The vertical maps are isomorphisms since the complements of U and $\pi^{-1}(U)$ have codimension at least 2 (for example, by [Har94, Proposition 1.11]). We conclude that the top map is an isomorphism, as required. $\square$

PROPOSITION 4.2. *Let $\pi: Y \rightarrow X$ be a proper morphism of varieties, where Y is smooth and X has rational singularities. Suppose that there is an open dense subset U of X such that $\pi^{-1}(U)$ is isomorphic (as a variety over U) to $U \times W$ for some irreducible projective variety W with rational singularities and satisfying $H^i(W, \mathcal{O}_W) = 0$ for $i > 0$. Then $R\pi_*(\mathcal{O}_Y) = \mathcal{O}_X$.*

Proof. Let $Z \subset Y \times W = Y \times_X (X \times W)$ be the closure of the graph of the isomorphism $\pi^{-1}(U) \rightarrow U \times W$, and let $\tilde{Z} \rightarrow Z$ be a resolution of singularities. Consider the diagram

$$\begin{array}{ccc} \tilde{Z} & \xrightarrow{\pi'} & X \times W \\ \rho' \downarrow & & \downarrow \rho \\ Y & \xrightarrow{\pi} & X. \end{array}$$

Since Z is the closure of the graph of an isomorphism of open sets, the projection maps $Z \rightarrow Y$ and $Z \rightarrow X \times W$ are birational. Thus π' and ρ' are birational; also, all maps above are proper. We now have

$$R\pi_*(\mathcal{O}_Y) = R\pi_*(R\rho'_*(\mathcal{O}_{\tilde{Z}})) = R\rho_*(R\pi'_*(\mathcal{O}_{\tilde{Z}})) = R\rho_*(\mathcal{O}_{X \times W}) = \mathcal{O}_X.$$

In the first step, we used that $R\rho'_*(\mathcal{O}_{\tilde{Z}}) = \mathcal{O}_Y$ since ρ' is a proper birational map of smooth varieties; in the second step, we used the commutativity of the diagram; in the third step we used that $R\pi'_*(\mathcal{O}_{\tilde{Z}}) = \mathcal{O}_{X \times W}$ since π' is a resolution of $X \times W$, which has rational singularities; and in the final step, we used the assumption on the cohomology of $\mathcal{O}_W$. $\square$

4.2 Some linear algebra

In this subsection, we work over an algebraically closed field $\mathbf{k}$ of arbitrary characteristic. Consider linear maps $f: V_0 \rightarrow V_1$ and $g: V_1 \rightarrow V_0$ such that V_0 and V_1 are finite-dimensional vector spaces. By picking bases, what form can we put the matrices in?

To answer this question, note that the tuples (V_0, V_1, f, g) form an abelian category $\mathcal{A}$. Concisely, we can regard $V_0 \oplus V_1$ as a $\mathbf{Z}/2$ -graded module over $\mathbf{k}[t]$, where t has degree 1; t acts on V_0 by f and on V_1 by g . We define some basic objects in this category.

- For $\lambda \in \mathbf{k}$, let $A_n(\lambda)$ be the object with $V_0 = V_1 = \mathbf{k}^n$ and where f is the identity matrix and g is a single Jordan block with eigenvalue λ .
- Let $A_n(\infty)$ be the object with $V_0 = V_1 = \mathbf{k}^n$ and where f is a single nilpotent Jordan block and g is the identity.

- Let B_n be the module $\mathbf{k}[t]/(t^{2n+1})$. This has basis $x_0, \dots, x_{2n}$, where x_i has parity i and $tx_i = x_{i+1}$ for $i < 2n$, and $tx_{2n} = 0$.

For an object M of $\mathcal{A}$, we let $M[1]$ be the object where the even and odd pieces of M are swapped. One easily verifies that $A_n(\lambda)[1]$ is isomorphic to $A_n(\lambda^{-1})$ for $\lambda \in \mathbf{k} \cup \{\infty\}$.

PROPOSITION 4.3. *The $A_n(\lambda)$ (for $\lambda \in \mathbf{k} \cup \{\infty\}$), B_n , and $B_n[1]$ are exactly the indecomposable objects of $\mathcal{A}$.*

Proof. Consider an object (V_0, V_1, f, g) . Then V_0 and V_1 are $\mathbf{k}[t^2]$ -modules, and f and g are maps of modules such that $fg = gf = t^2$. By the structure theorem for modules over a PID, V_0 decomposes as $V'_0 \oplus V''_0$, where t^2 is invertible on V'_0 and nilpotent on V''_0 ; of course, V_1 decomposes similarly. Since f and g are maps of modules, they respect these decompositions. Thus the whole object decomposes into two pieces, and it suffices to consider these pieces separately.

First suppose that t^2 is invertible. Since $fg = t^2$, it follows that $f: V_0 \rightarrow V_1$ is an isomorphism. We thus see that our object is isomorphic to the object $(V_1, V_1, \text{id}, fg)$, which is easily seen to decompose in terms of the $A_n(\lambda)$ with $\lambda \in \mathbf{k} \setminus \{0\}$.

Now suppose that t^2 is nilpotent. Then the usual proof of the structure theorem for modules over a PID applies, and we see that our module decomposes into cyclic modules, which are B_n , $B_n[1]$, $A_n(0)$, or $A_n(\infty)$. $\square$

REMARK 4.4. The category $\mathcal{A}$ is the representation category of the extended Dynkin quiver $\tilde{A}_1$. For more in this direction, see [DW17, § 7.1].

5. Grothendieck–Springer theory for the super Grassmannian

In this section, we study the Grothendieck–Springer theory associated with the super Grassmannian $\mathbf{Gr}_{r|s}(\mathbf{C}^{n|m})$ (see § 2.4). We prove a number of results about the various spaces. These results are used to compute the coherent cohomology of $\mathbf{Gr}_{r|s}(\mathbf{C}^{n|m})$ in the next section. This section takes place entirely in the world of ordinary (non-super) mathematics.

5.1 Statement of results

We introduce a number of objects:

- Let V_0 and V_1 be complex vector spaces of dimensions n and m .
- Let r and s be non-negative integers with $r \geq s$. Put

$$\delta = \begin{cases} m - n + r - s & \text{if } r - s > n - m, \\ 0 & \text{if } n - m \geq r - s. \end{cases}$$

Note that $m - \delta \leq \min(n, m)$.

- Put $W_0 = \text{Hom}(V_0, V_1)$ and $W_1 = \text{Hom}(V_1, V_0)$ and $W = W_0 \times W_1$. We regard these as affine varieties.
- Let $Z_0 \subset W_0$ be the determinantal variety consisting of linear maps of rank at most $m - \delta$. Let $Z_1 = W_1$, and put $Z = Z_0 \times Z_1$. (Note: when $\delta = 0$, we have $Z_0 = W_0$. This case is still interesting, though somewhat simpler.)
- Letting $f: V_0 \otimes \mathcal{O}_Z \rightarrow V_1 \otimes \mathcal{O}_Z$ and $g: V_1 \otimes \mathcal{O}_Z \rightarrow V_0 \otimes \mathcal{O}_Z$ denote the universal linear maps, let $\chi(u) \in \mathcal{O}_Z[u]$ be the characteristic polynomial of fg , and let $\bar{\chi}(u) = u^{-\delta}\chi(u)$. Since fg has rank at most $m - \delta$, it follows that $\bar{\chi}(u)$ is a polynomial.

- Let $\tilde{Z}$ be the affine scheme whose coordinate ring is the $(s, m - \delta - s)$ -factorization ring for $\bar{\chi}(u)$ introduced in § 3.7. Thus for a $\mathbf{C}$ -algebra T , a T -point of $\tilde{Z}$ is a triple (f, g, p) , where (f, g) is a T -point of Z and $p = p(u)$ is a degree s monic polynomial over T that divides $\bar{\chi}(u) \in T[u]$. There is a natural map $\tilde{Z} \rightarrow Z$ given by forgetting p .
- Let Y be the scheme defined as follows: a T -point of Y is a tuple (f, g, R_0, R_1) , where
 - * $R_0 \subset (V_0)_T$ is a T -submodule that is locally a rank r summand,
 - * $R_1 \subset (V_1)_T$ is a T -submodule that is locally a rank s summand,
 - * $f: (V_0)_T \rightarrow (V_1)_T$ is a map of T -modules such that $f(R_0) \subseteq R_1$,
 - * $g: (V_1)_T \rightarrow (V_0)_T$ is a map of T -modules such that $g(R_1) \subseteq R_0$.
 One easily sees that Y is the total space of a vector bundle over $\mathbf{Gr}_r(V_0) \times \mathbf{Gr}_s(V_1)$ and is thus smooth and irreducible.
- Let $\pi: Y \rightarrow \tilde{Z}$ be the map taking (f, g, R_0, R_1) to (f, g, p) , where p is the characteristic polynomial of fg on R_1 . We show that this is well defined in Corollary 5.6.

This setup turns out to be the Grothendieck–Springer theory associated with the super Grassmannian, as explained in § 6.2. The purpose of this section is to study the above situation, especially the varieties Y and $\tilde{Z}$. Our main results are summarized in the following theorem.

THEOREM 5.1. *We have the following:*

- (a) *The scheme $\tilde{Z}$ is integral and has rational singularities (and is thus normal and Cohen–Macaulay).*
- (b) *The map $\tilde{Z} \rightarrow Z$ is finite and flat; in fact, $\mathcal{O}_{\tilde{Z}}$ is a free $\mathcal{O}_Z$ -module of rank $\binom{m-\delta}{s}$.*
- (c) *The graded ring $\mathcal{O}_{\tilde{Z}} \otimes_{\mathcal{O}_Z} \mathbf{C}$ is isomorphic to $H_{\text{sing}}^*(\mathbf{Gr}_s(\mathbf{C}^{m-\delta}), \mathbf{C})$.*
- (d) *The map $\pi: Y \rightarrow \tilde{Z}$ is projective and satisfies $\pi_*(\mathcal{O}_Y) = \mathcal{O}_{\tilde{Z}}$ and $R^i \pi_*(\mathcal{O}_Y) = 0$ for $i > 0$. If $\delta > 0$ or $r = s$, then π is birational. If $\delta = 0$, then there is an open dense subset U of $\tilde{Z}$ such that $\pi^{-1}(U)$ is isomorphic to $U \times \mathbf{Gr}_{r-s}(\mathbf{C}^{n-m})$.*

REMARK 5.2. The proofs for the various parts are spread out:

- (a) The integrality is the content of § 5.4, and the fact that $\tilde{Z}$ has rational singularities is proven in Proposition 5.15.
- (b),(c) These are deduced in § 5.3.
- (d) The first claim is Proposition 5.17, the second claim is Proposition 5.13, and the third claim is Proposition 5.16.

REMARK 5.3. We make a clarifying remark about gradings. As W is a vector space, it carries a natural $\mathbf{G}_m$ -action via usual scalar multiplication, and this action induces all the gradings. In terms of rings, we have

$$\mathcal{O}_{W_0} = \text{Sym}(V_0 \otimes V_1^*), \quad \mathcal{O}_{W_1} = \text{Sym}(V_1 \otimes V_0^*), \quad \mathcal{O}_W = \mathcal{O}_{W_0} \otimes \mathcal{O}_{W_1},$$

and the elements of $V_0 \otimes V_1^*$ and $V_1 \otimes V_0^*$ are given degree 1. The space Y is a subbundle of $X_{\text{ord}} \times W$, where $X_{\text{ord}} = \mathbf{Gr}_r(V_0) \times \mathbf{Gr}_s(V_1)$; again, the $\mathbf{G}_m$ -action on W induces the grading on $\mathcal{O}_{\tilde{Z}} = \Gamma(Y, \mathcal{O}_Y)$. Precisely, writing $Y = \text{Sym}(\eta)$, where η is a quotient of $\mathcal{O}_{X_{\text{ord}}} \otimes W^*$, the degree d piece of $\mathcal{O}_{\tilde{Z}}$ is $\Gamma(X_{\text{ord}}, \text{Sym}^d(\eta))$. We note that the gradings we use do not necessarily coincide with the gradings induced by the central tori in $\mathbf{GL}(V_0)$ and $\mathbf{GL}(V_1)$.

5.2 Some auxiliary spaces

To prove Theorem 5.1, we will use a few auxiliary spaces, which we now introduce. Conceptually, these spaces come from the Grothendieck–Springer theory associated with a flag supervariety.

- Let $\tilde{\mathcal{Z}}$ be the affine scheme whose coordinate ring is the splitting ring for $\bar{\chi}(u)$ introduced in § 3.1.
 - Let $\mathcal{Y}$ be the scheme defined as follows: a T -point of $\mathcal{Y}$ is a tuple $(f, g, F_\bullet, G_\bullet)$, where
 - * $F_{r-s} \subset F_{r-s+1} \subset \cdots \subset F_{r-s+m-\delta} \subset (V_0)_T$ is a flag of T -submodules which are locally summands with ranks prescribed by the subscript,
 - * $0 = G_0 \subset G_1 \subset \cdots \subset G_{m-\delta} \subset (V_1)_T$ is a flag of T -submodules which are locally summands with ranks prescribed by the subscript,
 - * $f: (V_0)_T \rightarrow (V_1)_T$ is a map of T -modules such that $f(F_{i+r-s}) \subseteq G_i$ for all $i \geq 0$,
 - * $g: (V_1)_T \rightarrow (V_0)_T$ is a map of T -modules such that $g(G_i) \subseteq F_{i+r-s}$ for all $i \geq 0$.
- In fact, $\mathcal{Y}$ is the total space of a vector bundle over a product of partial flag varieties $\mathbf{Fl}(r-s, \dots, r-s+m-\delta; V_0) \times \mathbf{Fl}(1, 2, \dots, m-\delta; V_1)$.
- Let $\rho: \mathcal{Y} \rightarrow \tilde{\mathcal{Z}}$ be the map taking $(f, g, F_\bullet, G_\bullet)$ to $(f, g, \prod_{i=1}^{m-\delta}(u - \lambda_i))$, where λ_i is the eigenvalue of fg on G_i/G_{i-1} . We show that this is well defined in Proposition 5.5.

We prove the following analog of Theorem 5.1.

THEOREM 5.4. *We have the following:*

- (a) *The scheme $\tilde{\mathcal{Z}}$ is integral and has rational singularities (and is thus normal and Cohen–Macaulay).*
- (b) *The map $\tilde{\mathcal{Z}} \rightarrow Z$ is finite and flat; in fact, $\mathcal{O}_{\tilde{\mathcal{Z}}}$ is a free $\mathcal{O}_Z$ -module of rank $(m-\delta)!$.*
- (c) *The graded ring $\mathcal{O}_{\tilde{\mathcal{Z}}} \otimes_{\mathcal{O}_Z} \mathbf{C}$ is isomorphic to $H_{\text{sing}}^*(\mathbf{Fl}(\mathbf{C}^{m-\delta}), \mathbf{C})$.*
- (d) *The map $\rho: \mathcal{Y} \rightarrow \tilde{\mathcal{Z}}$ is projective and satisfies $\rho_*(\mathcal{O}_{\mathcal{Y}}) = \mathcal{O}_{\tilde{\mathcal{Z}}}$ and $R^i \rho_*(\mathcal{O}_{\mathcal{Y}}) = 0$ for $i > 0$. If $\delta > 0$ or $r = s$, then ρ is birational. If $\delta = 0$, then there is an open dense subset U of $\tilde{\mathcal{Z}}$ such that $\rho^{-1}(U)$ is isomorphic to $U \times \mathbf{Gr}_{r-s}(\mathbf{C}^{n-m})$.*

See Remark 5.2 for the locations of the proofs of the various statements (we handle this and the previous case in parallel).

5.3 First observations

It is well known that Z_0 has rational singularities [Wey03, Proposition 6.2.3] and is in particular normal and Cohen–Macaulay. Since Z_1 is an affine space, it follows that the same holds for $Z = Z_0 \times Z_1$. Theorem 5.1(b),(c) and Theorem 5.4(b),(c), as well as the fact that $\tilde{Z}$ and $\tilde{\mathcal{Z}}$ are Cohen–Macaulay, follow immediately from Propositions 3.1, 3.4, 3.11, and 3.13. The following proposition and corollary ensure that ρ and π are well defined.

PROPOSITION 5.5. *Suppose that $(f, g, F_\bullet, G_\bullet)$ is a T -point of $\mathcal{Y}$. Let λ_i be the eigenvalue of fg on G_i/G_{i-1} . Then $\prod_{i=1}^{m-\delta}(u - \lambda_i) = \bar{\chi}(u)$.*

Proof. If $\delta = 0$, then $G_{m-\delta} = V_1$ and $\bar{\chi}(u) = \chi(u)$, so there is nothing to prove. Otherwise, suppose $\delta = m - n + r - s > 0$. Then $F_{r-s+m-\delta} = V_0$, so that the image of f is contained in $G_{m-\delta}$. So fg induces the zero endomorphism on $V_1/G_{m-\delta}$, which implies that the characteristic polynomial of fg on $G_{m-\delta}$ is $\bar{\chi}(u)$, and which proves the result. $\square$

COROLLARY 5.6. *Suppose that (f, g, R_0, R_1) is a T -point of Y . Let p be the characteristic polynomial of fg on R_1 . Then $p(u)$ divides $\bar{\chi}(u)$.*

Proof. This is immediate from the fact that we have a surjective map $\mathcal{Y} \rightarrow Y$ given by

$$(f, g, F_\bullet, G_\bullet) \mapsto (f, g, F_r, G_s).$$

□

5.4 Integrality

We now show that $\tilde{Z}$ and $\tilde{\mathcal{Z}}$ are integral. Let $\Delta \in \mathcal{O}_Z$ be the discriminant of $\bar{\chi}$. We begin with the following.

PROPOSITION 5.7. *We have $\Delta \neq 0$, and so $\tilde{Z}$ and $\tilde{\mathcal{Z}}$ are reduced.*

Proof. We construct a $\mathbf{C}$ -point of Z where $\Delta \neq 0$. Let $x_1, \dots, x_n$ and $y_1, \dots, y_m$ be bases for V_0 and V_1 . Let $\lambda_1, \dots, \lambda_{m-\delta}$ be distinct complex numbers. Define $f: V_0 \rightarrow V_1$ by $f(x_i) = \lambda_i y_i$ for $1 \leq i \leq m - \delta$ and $f(x_i) = 0$ for $i > m - \delta$. Define $g: V_1 \rightarrow V_0$ by $g(x_i) = y_i$ for $1 \leq i \leq m - \delta$ and $g(y_i) = 0$ for $i > m - \delta$. Note that (f, g) defines a $\mathbf{C}$ -point of Z . The composition fg has characteristic polynomial $\chi(u) = u^\delta \prod_{i=1}^{m-\delta} (u - \lambda_i)$. Thus $\Delta \neq 0$ at (f, g) . Since Z is integral, it follows that Δ is not a zerodivisor, and so $\tilde{Z}$ is reduced (Proposition 3.1(e)), and so $\tilde{\mathcal{Z}}$ is as well by Proposition 3.11(e). □

To complete the proof of the integrality, it suffices to show that $\tilde{Z}$ and $\tilde{\mathcal{Z}}$ are irreducible. Since Y and $\mathcal{Y}$ are irreducible, this is a consequence of the following result.

PROPOSITION 5.8. *The maps $\rho: \mathcal{Y} \rightarrow \tilde{\mathcal{Z}}$ and $\pi: Y \rightarrow \tilde{Z}$ are surjective.*

Proof. Let $(f, g, \prod_{i=1}^{m-\delta} (u - \lambda_i))$ be a $\mathbf{C}$ -point of $\tilde{\mathcal{Z}}$. Recall the category $\mathcal{A}$ from § 4.2. Let M be the object (V_0, V_1, f, g) . We inductively construct a chain of subobjects $M^0 \subset \dots \subset M^{m-\delta}$ of M , with the following properties:

- (a) We have $\dim(M_0^i) = i + r - s$ and $\dim(M_1^i) = i$.
- (b) For $1 \leq i \leq m - \delta$, the eigenvalues of t^2 on M^i/M^{i-1} are λ_i .

To begin, we take M_0^0 to be a subspace of $\ker(f)$ of dimension $r - s$ (which exists since $f \in Z_0$) and $M_1^0 = 0$. Now suppose that we have defined M^{i-1} for some $1 \leq i \leq m - \delta$.

We claim that there is an $\mathcal{A}$ -subobject of M/M^{i-1} of dimension $1|1$ such that the eigenvalues of t^2 are both λ_i ; granted this, we take M^i/M^{i-1} to be this subobject. By considering the characteristic polynomial of $t^2 = fg$ on M_1 , we see that λ_i occurs as an eigenvalue of t^2 on M_1^i/M_1^{i-1} . If $\lambda_i \neq 0$, then the claim follows from the form of indecomposables given in § 4.2. Now suppose $\lambda_i = 0$. For an object N of $\mathcal{A}$, let $\Phi(N)$ be the generalized 0-eigenspace of t^2 , and let $\Psi(N)$ be the sum of the generalized λ -eigenspaces with $\lambda \neq 0$. We have $\dim(\Psi(N)_0) = \dim(\Psi(N)_1)$. Let k be the number of λ_i that are zero. Then

$$\dim(\Phi(M)_1) = \delta + k, \quad \dim(\Psi(M)_1) = m - (\delta + k), \quad \dim(\Phi(M)_0) = n - m + \delta + k.$$

We thus find

$$\dim(\Phi(M/M^0)_0) = n - m + s - r + \delta + k \geq k.$$

Now, as $\lambda_i = 0$, it follows that fewer than k of $\lambda_1, \dots, \lambda_{i-1}$ are 0, and so $\dim(\Phi(M/M^{i-1})_0) > 0$. We thus see that $\Phi(M/M^{i-1})$ has non-zero even and odd parts and thus (by the classification of indecomposables) has a subobject of dimension $1|1$.

We now define a $\mathbf{C}$ -point $(f, g, F_\bullet, G_\bullet)$ of $\mathcal{Y}$ by taking $F_{r-s+i} = M_0^i$ and $G_i = M_1^i$. It is clear that this is indeed a $\mathbf{C}$ -point of $\mathcal{Y}$ and lifts the point of $\tilde{\mathcal{Z}}$ under consideration. Thus ρ is

surjective. We have a commutative diagram

$$\begin{array}{ccc} \mathcal{Y} & \longrightarrow & Y \\ \rho \downarrow & & \downarrow \pi \\ \tilde{\mathcal{Z}} & \longrightarrow & \tilde{Z}, \end{array}$$

where the top map takes $(f, g, F_\bullet, G_\bullet)$ to (f, g, F_r, G_s) and the bottom map is the natural one. Since the bottom map is clearly surjective, so is π . $\square$

5.5 Normality

We now aim to show that $\tilde{Z}$ and $\tilde{\mathcal{Z}}$ are normal. We will use the criterion in Proposition 3.10. We therefore begin by studying the geometry of $V(\Delta)$ and $V(\Delta, \partial\Delta)$.

PROPOSITION 5.9. *Let (f, g) be a $\mathbf{C}$ -point of Z . We can choose bases of V_0 and V_1 such that the matrices for f and g have the form*

$$\begin{pmatrix} 0 & * \\ 0 & A \end{pmatrix}, \quad \begin{pmatrix} * & * \\ 0 & B \end{pmatrix},$$

where A and B are upper-triangular square matrices with $m - \delta$ rows and columns (this determines the sizes of the other blocks). Moreover, if $(f, g) \in V(\Delta)$, then one can also assume that the first two diagonal entries of AB are equal.

Proof. Since f has rank at most $m - \delta$, its kernel has dimension at least $n - m + \delta$; similarly, since fg has rank at most $m - \delta$, its kernel has dimension at least δ . Let $U_1 \subset \ker(fg)$ be a subspace of dimension δ . Then $g(U_1)$ is a subspace of $\ker(f)$ of dimension at most δ . Let $U_0 \subset \ker(f)$ be a subspace of dimension $n - m + \delta$ containing $g(U_1)$. Now, pick a basis $x_1, \dots, x_n$ of V_0 such that $x_1, \dots, x_{n-m+\delta}$ is a basis of U_0 , and pick a basis $y_1, \dots, y_m$ of V_1 such that $y_1, \dots, y_\delta$ is a basis for U_1 . In these bases, the matrices for f and g have the stated form, except that A and B may not be upper-triangular. Let U'_0 be the span of $x_{n-m+\delta+1}, \dots, x_n$, and let U'_1 be the span of $y_{\delta+1}, \dots, y_m$. We can then regard (A, B, U'_0, U'_1) as an object of the category considered in § 4.2. By decomposing into indecomposables and choosing bases for these indecomposables, we can make A and B upper-triangular. (The one subtlety here is that the matrices for B_n and B'_n are not square. However, since $\dim(U'_0) = \dim(U'_1)$, whenever we have a B_n , we can pair it with some B'_m , and the matrices for $B_n \oplus B'_m$ are square and can be taken to be upper-triangular.)

Finally, suppose $(f, g) \in V(\Delta)$. If $\bar{\chi}$ has a non-zero repeated root λ , then by taking the $A_n(\lambda)$ first in the decomposition, we see that the first two diagonal entries of AB are λ . Otherwise, 0 is a repeated root of $\bar{\chi}$, and by putting the $A_n(0)$, $A_n(\infty)$, B_n , and B'_n first in the decomposition, the first two diagonal entries of AB are 0. $\square$

PROPOSITION 5.10. *The closed set $V(\Delta) \subset Z$ is irreducible; in fact, any two points of $V(\Delta)$ can be joined by an irreducible rational curve in $V(\Delta)$.*

Proof. Fix bases for V_0 and V_1 , and let (f_i, g_i) for $i = 1, 2$ be two points of $V(\Delta)$. Applying Proposition 5.9, we can find $\gamma_i \in \mathbf{GL}(V_0) \times \mathbf{GL}(V_1)$ such that

$$\gamma_i f_i = \begin{pmatrix} 0 & C_i \\ 0 & A_i \end{pmatrix}, \quad \gamma_i g_i = \begin{pmatrix} D_i & E_i \\ 0 & B_i \end{pmatrix},$$

where A_i and B_i are upper-triangular square matrices of size $m - \delta$ and the first two diagonal entries of $A_i B_i$ are equal. We define matrices $A(t), \dots, E(t)$ with polynomial entries such that the following hold:

- At $t = 0$, they coincide with $A_1, \dots, E_1$.
- At $t = 1$, they coincide with $A_2, \dots, E_2$.
- The matrices $A(t)$ and $B(t)$ are upper-triangular.
- The first two diagonal entries of $A(t)B(t)$ agree.

We take $C(t) = tC_1 + (1-t)C_2$. We use the same formula everywhere else, except for the first two diagonal entries of $A(t)$ and $B(t)$. Consider the subvariety of $\mathbf{A}^4$ defined by $\alpha_1\alpha_2 = \beta_1\beta_2$. Any two points on this variety can be joined by a map from $\text{Spec}(\mathbf{C}[t])$, as it is the affine cone over $\mathbf{P}^1 \times \mathbf{P}^1$. We take $(A_{1,1}(t), A_{2,2}(t), B_{1,1}(t), B_{2,2}(t))$ to be such a curve joining $(A_{i,1,1}, A_{i,2,2}, B_{i,1,1}, B_{i,2,2})$.

Now, define

$$F(t) = \begin{pmatrix} 0 & C(t) \\ 0 & A(t) \end{pmatrix}, \quad G(t) = \begin{pmatrix} D(t) & E(t) \\ 0 & B(t) \end{pmatrix}.$$

Since $\ker(F(t))$ contains the first $n - m + \delta$ basis vectors, the rank of $F(t)$ is $\leq m - \delta$. Thus $(F(t), G(t))$ defines a point of Z . The polynomial $\bar{\chi}_{F(t), G(t)}$ is the characteristic polynomial of $A(t)B(t)$, which has a repeated root since the first two diagonal entries coincide. Thus $(F(t), G(t))$ is a point of $V(\Delta)$.

Finally, let $\gamma: \text{Spec}(\mathbf{C}[t]) \rightarrow \mathbf{GL}(V_0) \times \mathbf{GL}(V_1)$ be a curve such that $\gamma(0) = \gamma_1$ and $\gamma(1) = \gamma_2$. Then $\gamma(F(t), G(t))$ is a curve in $V(\Delta)$ joining (f_1, g_1) to (f_2, g_2) . This completes the proof. $\square$

PROPOSITION 5.11. *The schemes $\tilde{Z}$ and $\tilde{Z}$ are normal.*

Proof. It suffices to show that $\tilde{Z}$ is normal (due to Proposition 3.11(e)). If $m - \delta \leq 1$, then $\deg(\bar{\chi}) \leq 1$ and $\tilde{Z} = Z$ is normal. We thus assume $m - \delta \geq 2$ in what follows.

We verify the conditions of Proposition 3.10. We already know that $\mathcal{O}_Z$ is normal and that Δ is not a zerodivisor (Proposition 5.7). It thus suffices to show that $V(\Delta, \partial\Delta)$ has codimension at least 2 in Z . Since Z is normal, its singular locus has codimension at least 2. It is thus enough to show that $Z_{\text{reg}} \cap V(\Delta, \partial\Delta)$ has codimension at least 2 in Z_{reg} , where Z_{reg} is the regular locus (we will use below that Z_{reg} is the set of matrices with maximal possible rank). The set $Z_{\text{reg}} \cap V(\Delta, \partial\Delta)$ is closed in Z_{reg} (see the first paragraph of § 3.6) and contained in $V(\Delta)$. Since $V(\Delta)$ is irreducible of codimension 1, it thus suffices to show that $Z_{\text{reg}} \cap V(\Delta, \partial\Delta)$ is a proper subset of $Z_{\text{reg}} \cap V(\Delta)$. We do this by writing down a point in $Z_{\text{reg}} \cap V(\Delta)$ that does not belong to $V(\Delta, \partial\Delta)$.

Pick bases for V_0 and V_1 . We define a $\mathbf{C}[\epsilon]/(\epsilon^2)$ point of Z by

$$f = \begin{pmatrix} 0 & 0 \\ 0 & A \end{pmatrix}, \quad g = \begin{pmatrix} 0 & 0 \\ 0 & I \end{pmatrix},$$

where A and I are square matrices of size $m - \delta$, I is the identity matrix, and

$$A = \begin{pmatrix} \lambda & 1 & & \\ \epsilon & \lambda & & \\ & & & D \end{pmatrix},$$

where $\lambda \in \mathbf{C} \setminus \{0\}$ and D is a diagonal matrix with distinct non-zero complex entries that are not λ . Let $x = (\bar{f}, \bar{g})$ be the $\mathbf{C}$ -point of Z obtained by putting $\epsilon = 0$. It is clear that x belongs to $V(\Delta)$ since $\bar{f}\bar{g}$ has a repeated eigenvalue. Since $\bar{f}$ has maximal possible rank, it is a smooth point of Z_0 , and so $x \in Z_{\text{reg}}$. The value of Δ on (f, g) is a non-zero multiple of ϵ : the key point is that the discriminant of the characteristic polynomial of the top 2×2 block of A is 4ϵ . This shows that $d\Delta$ is non-zero at (f, g) , and so $x \notin V(\Delta, \partial\Delta)$. $\square$

5.6 Rational singularities

We now show that $\tilde{Z}$ and $\tilde{\mathcal{Z}}$ have rational singularities. To this end, we introduce the following closed subsets of Z :

- Let D_1 be the locus of points such that 0 is a root of $\bar{\chi}(u)$.
- Let D_2 be the locus such that $\bar{\chi}(u)$ has a repeated root.
- Let $D_3 \subset D_2$ be the locus where there is
 - * a triple root, or
 - * two repeated roots, or
 - * a unique repeated root, but the corresponding Jordan block of fg is a scalar.

This set is closed, as we now explain. The first two conditions define a closed set. Letting U be the complement, it suffices to show that $D_3 \cap U$ is closed in U . Let $V \subset U \times \mathbf{A}^1$ be the set of pairs (f, g, λ) where λ is a root of $\bar{\chi}$ and its derivative. The map $V \rightarrow U$ is finite and injective with image $V(\Delta) \cap U$. Let $V_1 \subset V$ be the closed set where $fg - \lambda$ has nullity at least 2. Then $D_3 \cap U$ is the image of V_1 in U and thus closed.

- Let $D_4 = D_1 \cap D_2$, and let $D_5 = D_1 \cup D_3$.

Put $U_i = Z \setminus D_i$. For any space X over Z , we let $D_i(X)$ and $U_i(X)$ denote the inverse images of D_i and U_i , respectively, in X .

PROPOSITION 5.12. *The sets $D_3(\mathcal{Y})$ and $D_4(\mathcal{Y})$ have codimension at least 2 in $\mathcal{Y}$.*

Proof. By equivariance, the restrictions of $D_3(\mathcal{Y})$ or $D_4(\mathcal{Y})$ to any fiber over $\mathbf{Fl}(r - s, \dots, r - s + m - \delta; V_0) \times \mathbf{Fl}(1, \dots, m - \delta; V_1)$ is isomorphic to any other. So it suffices to show that in a given fiber of the vector bundle, these restrictions have codimension at least 2. In that case, by picking bases of V_0 and V_1 adapted to the particular pair of flags, f and g have the following form:

- The first $r - s$ columns of f are 0.
- The next $m - \delta$ columns of f are upper-triangular.
- The remaining columns of f (if they exist) are arbitrary.
- The first $r - s$ rows of g are arbitrary.
- The next $m - \delta$ rows of g are upper-triangular.
- The remaining rows of g (if they exist) are 0.

So fg is determined by the contents of the upper-triangular $m - \delta$ columns of f and the upper-triangular $m - \delta$ rows of g . Let $x_1, \dots, x_{m-\delta}$ and $y_1, \dots, y_{m-\delta}$ be the corresponding diagonal entries of f and g , respectively.

Hence over this fiber, the condition that a root is repeated has one component for each equation $x_i y_i = x_j y_j$ for $i < j$. Then $D_4(\mathcal{Y})$ is cut out in this component by $\prod_i x_i y_i$ and hence has codimension 2. Having a triple root or two repeated roots corresponds to imposing two equations $x_i y_i = x_j y_j$ and $x_k y_k = x_\ell y_\ell$, so has codimension 2. If there is a repeated root which is unique, the condition that the Jordan block is scalar is a non-trivial condition on an irreducible component and hence also has codimension at least 2. $\square$

PROPOSITION 5.13. *Assume $r = s$ or $\delta > 0$. Then $\rho: U_5(\mathcal{Y}) \rightarrow U_5(\tilde{\mathcal{Z}})$ is an isomorphism, and so ρ is birational. Similarly, $\pi: U_5(Y) \rightarrow U_5(\tilde{Z})$ is an isomorphism, and so π is birational.*

Proof. We first prove the statement about ρ . Since $\tilde{\mathcal{Z}}$ is normal and ρ is projective, it suffices to show that ρ induces an isomorphism on $\mathbf{C}$ -points on these two sets. By Proposition 5.8, we know that ρ is surjective, so it suffices to prove injectivity.

First suppose $r = s$. Consider a point $(f, g, \prod_{i=1}^{m-\delta}(u - \lambda_i))$ in $U_5(\tilde{\mathcal{Z}})$, and let $(f, g, F_\bullet, G_\bullet)$ be an inverse image under ρ . If the λ_i are distinct, then G_i must be the span of the fg -eigenspaces for $\lambda_1, \dots, \lambda_i$ and F_i must be the span of the gf -eigenspaces for $\lambda_1, \dots, \lambda_i$. Now suppose that there is a repeated root. Since our point belongs to $U_5(\tilde{\mathcal{Z}})$, there is only one repeated root, and the corresponding Jordan block for fg is non-scalar. The spaces F_i and G_i are again uniquely determined: for the repeated eigenvalue, we use the eigenvector first and the generalized eigenvector second. This shows that there is a unique inverse image under ρ .

Now suppose instead that $\delta > 0$. As before, there is a unique choice of the flag G_i . Also by our definition of U_1 , we have $\text{rank}(fg) = m - \delta = \text{rank}(gf)$ and hence $\dim \ker(gf) = r - s$. Hence F_i must be the span of $\ker(gf)$ and the eigenspaces for $\lambda_1, \dots, \lambda_i$.

The claim about π follows from similar reasoning. $\square$

COROLLARY 5.14. *Assume $r = s$ or $\delta > 0$. Then $U_2(\tilde{\mathcal{Z}})$ and $U_5(\tilde{\mathcal{Z}})$ have rational singularities.*

Proof. By Proposition 5.13, the variety $U_5(\tilde{\mathcal{Z}})$ is smooth and hence has rational singularities. Next, $\tilde{\mathcal{Z}} \rightarrow Z$ is étale over U_2 , and since Z has rational singularities, the same is true for $U_2(\tilde{\mathcal{Z}})$. $\square$

PROPOSITION 5.15. *The varieties $\tilde{\mathcal{Z}}$ and $\tilde{Z}$ have rational singularities.*

Proof. First suppose $\delta > 0$ or $r = s$. By Corollary 5.14, the scheme $\tilde{\mathcal{Z}}$ has rational singularities on $U_2(\tilde{\mathcal{Z}}) \cup U_5(\tilde{\mathcal{Z}})$. The complement is $D_3(\tilde{\mathcal{Z}}) \cap D_4(\tilde{\mathcal{Z}})$, and the inverse image of this set under ρ has codimension at least 2 by Proposition 5.12. Hence $\tilde{\mathcal{Z}}$ has rational singularities by Proposition 4.1. Next if $\delta = 0$ and $r \neq s$, the variety $\tilde{\mathcal{Z}}$ is the same as the variety $\tilde{\mathcal{Z}}$ with (r, s) changed to (s, s) , and thus has rational singularities as well. Finally, $\tilde{Z}$ is a quotient of $\tilde{\mathcal{Z}}$ by a finite group, so has rational singularities by [KM98, Proposition 5.13]. $\square$

5.7 Cohomology of Y and $\mathcal{Y}$

We now finish off the proofs by computing the cohomology of $\mathcal{O}_Y$ and $\mathcal{O}_{\mathcal{Y}}$. The following result is required to treat the $\delta = 0$ case and is part of Theorems 5.1(d) and 5.4(d). Recall that $\delta = 0$ is equivalent to the condition $n - m \geq r - s$.

PROPOSITION 5.16. *Suppose $\delta = 0$.*

- (a) *Let $U \subset Z$ be the locus where $fg: V_1 \rightarrow V_1$ is an isomorphism, $\Delta \neq 0$, and $\bar{\chi}(0) \neq 0$. Then U is an open dense subset of Z .*
- (b) *There is a vector bundle $\mathcal{E}$ on U of rank $n - m$ whose fiber at (f, g) is the kernel of f .*

Let U' be a dense open subset of U where $\mathcal{E}$ is trivial, and let $\tilde{U}$ and $\tilde{\mathcal{U}}$ be the inverse images of U' in $\tilde{\mathcal{Z}}$ and $\tilde{Z}$. Then

- (c) $\pi^{-1}(\tilde{U}) \cong \mathbf{Gr}_{r-s}(\mathbf{C}^{n-m}) \times \tilde{U}$, and
- (d) $\rho^{-1}(\tilde{\mathcal{U}}) \cong \mathbf{Gr}_{r-s}(\mathbf{C}^{n-m}) \times \tilde{\mathcal{U}}$.

Proof. (a) It is clear that U is open. We show that it is non-empty. Since $\delta = 0$, we have $n \geq m$. Let $x_1, \dots, x_n$ be a basis of V_0 and $y_1, \dots, y_m$ a basis of V_1 . Define $f: V_0 \rightarrow V_1$ by $f(x_i) = y_i$ for

$1 \leq i \leq m$ and $f(x_i) = 0$ for $i > m$. Define $g: V_1 \rightarrow V_0$ by $g(y_i) = \lambda_i x_i$, where $\lambda_1, \dots, \lambda_m$ are distinct non-zero complex numbers. Then (f, g) belongs to U .

(b) If $(f, g) \in U$, then f is surjective and $\ker(f)$ has dimension $n - m$, so we can take $\mathcal{E}$ to be the kernel of the generic map $f: V_0 \otimes \mathcal{O}_U \rightarrow V_1 \otimes \mathcal{O}_U$.

(c) Suppose that (f, g, p) is a point in $\tilde{U}$. Consider a point (f, g, R_0, R_1) in Y above (f, g, p) . Then R_1 is uniquely determined: it is the sum of the eigenspaces of fg corresponding to eigenvalues that are roots of p . Since fg is an isomorphism, g is injective and its image is linearly disjoint from $\ker(f)$. Thus $R_0 = g(R_1) \oplus K$, where K is a subspace of $\ker(f)$ of dimension $r - s$. Moreover, any choice of K leads to a point above (f, g, p) . We thus see that the fiber above (f, g, p) is $\mathbf{Gr}_{r-s}(\mathbf{C}^{n-m})$. The isomorphism $\pi^{-1}(\tilde{U}) = \mathbf{Gr}_{r-s}(\mathbf{C}^{n-m}) \times \tilde{U}$ comes from combining this analysis with a choice of trivialization of $\mathcal{E}$ over U' .

(d) This is similar to part (c): the flag $G_\bullet$ is uniquely determined, F_{r-s} is an $r - s$ dimensional subspace of $\ker(f)$, and $F_{r-s+i} = g(G_i) + F_{r-s}$ for $1 \leq i \leq m - \delta$. $\square$

PROPOSITION 5.17. We have $R\pi_*(\mathcal{O}_Y) = \mathcal{O}_{\tilde{Z}}$ and $R\rho_*(\mathcal{O}_Y) = \mathcal{O}_{\tilde{Z}}$.

Proof. If $\delta > 0$ or $r = s$, then π and ρ are birational (Proposition 5.13), and the result follows from generalities on rational singularities (see § 4.1). For $\delta = 0$, the result follows from Propositions 4.2 and 5.16. $\square$

6. Cohomology of the super Grassmannian

6.1 Statement of results

We use notation as in § 5.1. Additionally, we introduce the following notation.

- Let V be the super vector space $V_0 \oplus V_1$.
- Let X be the super Grassmannian $\mathbf{Gr}_{r|s}(V)$ (see § 6.2 for background).
- Let $\mathbf{G}$ be the super group $\mathbf{GL}(V)$, and let $\mathbf{G}_{\text{ord}} = \mathbf{GL}(V_0) \times \mathbf{GL}(V_1)$ be the underlying ordinary group.
- Let $A = H_{\text{sing}}^*(\mathbf{Gr}_s(\mathbf{C}^{m-\delta}), \mathbf{C})$, regarded as a graded $\mathbf{C}$ -algebra.
- Let $S = \text{Sym}(W^*)$ be the coordinate ring of W .

The main result of this section is Theorem 1.2, which computes the cohomology of $\mathcal{O}_X$. We restate the theorem here in our current notation:

THEOREM 6.1. We have the following:

- (a) We have a natural isomorphism $H^*(X, \mathcal{O}_X)^{\mathbf{G}} = A$ of graded algebras.
- (b) There is a canonical graded $\mathbf{G}$ -subrepresentation E of $H^*(X, \mathcal{O}_X)$ such that the natural map $A \otimes E \rightarrow H^*(X, \mathcal{O}_X)$ is an isomorphism.
- (c) We have a canonical isomorphism of $\mathbf{G}_{\text{ord}}$ -representations

$$E^i = \bigoplus_{p \geq 0} \text{Tor}_p^S(\mathcal{O}_Z, \mathbf{C})_{i+p}.$$

We note that if $\delta = 0$, then $Z = W$, and so $E^i = 0$ for $i > 0$ and $E^0 = \mathbf{C}$. Moreover, $\mathbf{G}_{\text{ord}}$ acts trivially on E^0 , and so $\mathbf{G}$ does as well. Thus, in this case, we have $H^*(X, \mathcal{O}_X) = A$ with trivial $\mathbf{G}$ -action, as in Theorem 1.2(a).

A corollary of the theorem is that the $\mathbf{G}_{\text{ord}}$ -action on the linear strands of the resolution of $\mathcal{O}_Z$ extends to an action of $\mathbf{G}$. This result was first proved by Pragacz–Weyman [PW85]. We discuss this in more detail in § 6.5.

6.2 Grothendieck–Springer theory

We now give some background on the super Grassmannian and connect it to varieties studied in § 5. While the super Grassmannian can be expressed in the form $\mathbf{G}/\mathbf{P}$ for an appropriate parabolic subgroup $\mathbf{P}$ of $\mathbf{G}$, we use an alternative approach here to be more direct. We refer to [Man97, Chapter 4, § 3] for general background on the super Grassmannian.

The super Grassmannian $X = \mathbf{Gr}_{r|s}(V)$ is the super scheme representing the functor that attaches to a super algebra T the set of T -submodules of $T \otimes V$ that are locally summands of rank $r|s$. It is not difficult to see that this functor is indeed representable. One can show that X is smooth using the criterion for formal smoothness in the super setting; this is also proved in [Man97]. The super scheme X is smooth of dimension $d_0|d_1$, where

$$d_0 = r(n - r) + s(m - s), \quad d_1 = r(m - s) + s(n - r).$$

Let

$$0 \rightarrow \mathcal{R} \rightarrow \mathcal{O}_X \otimes V \rightarrow \mathcal{Q} \rightarrow 0$$

be the tautological sequence on X : given a superalgebra T , a T -valued point of $\mathcal{R}$ is a pair (W, w) , where W is a T -submodule of $T \otimes V$ which is locally a summand of rank $r|s$ (that is, a T -valued point of X), together with a vector $w \in W$. Informally, the fiber of $\mathcal{R}$ over a submodule is the submodule itself. In particular, $\mathcal{R}$ is a vector bundle of rank $r|s$. Also, let X_0 be the ordinary scheme $\mathbf{Gr}_r(V_0)$, and let

$$0 \rightarrow \mathcal{R}_0 \rightarrow \mathcal{O}_{X_0} \otimes V_0 \rightarrow \mathcal{Q}_0 \rightarrow 0$$

be its tautological sequence (defined in a way similar to the above). Similarly, define $X_1 = \mathbf{Gr}_s(V_1)$, and let

$$0 \rightarrow \mathcal{R}_1 \rightarrow \mathcal{O}_{X_1} \otimes V_1 \rightarrow \mathcal{Q}_1 \rightarrow 0$$

be its tautological sequence. Restricting the functor of points of X to ordinary algebras, one sees that $X_{\text{ord}} = X_0 \times X_1$. One also finds $\mathcal{Q}_{\text{ord}} = \mathcal{Q}_0 \oplus \mathcal{Q}_1$, and similarly $\mathcal{R}_{\text{ord}} = \mathcal{R}_0 \oplus \mathcal{R}_1$. (We simply write $\mathcal{Q}_0$ for the pullback of $\mathcal{Q}_0$ to $X_0 \times X_1$, and similarly in other cases.)

We now determine $\text{gr}(\mathcal{O}_X)$. The result of the calculation below can be found in [Man97], but we prefer to give a short self-contained explanation. We begin with the following observation.

LEMMA 6.2. *Let T be a super algebra, let $J = J_T$, and let*

$$0 \rightarrow M \rightarrow T \otimes V \rightarrow N \rightarrow 0$$

be an exact sequence of T -modules. Then there exists a unique map of T/J -modules

$$\varphi: M/JM \rightarrow JN/J^2N$$

satisfying the following condition: if x is a homogeneous element of M of degree d and $x = y_0 + y_1$ with $y_i \in T_{d+i} \otimes V_i$, then $\varphi(\bar{x}) = \bar{y}_{d+1}$, where the bar denotes the image in the quotient module.

Proof. Define $\tilde{\varphi}: M \rightarrow J \otimes V$ as follows: given $x \in M_d$, write $x = y_0 + y_1$ with $y_i \in T_{i+d} \otimes V_i$, and put $\tilde{\varphi}(x) = y_{d+1}$. Note that $y_{d+1} \in T_1 \otimes V_{d+1}$ does indeed belong to $J \otimes V$. One easily sees that $\tilde{\varphi}$ is a map of T_0 -modules. Now, suppose that $x = y_0 + y_1$ is as above and $a \in T_1$. Then $ax = ay_0 + ay_1$ has degree $d + 1$, and so $\tilde{\varphi}(ax) = ay_d = ax - ay_{d+1}$ belongs to $JM + J^2 \otimes V$. It

follows that $\tilde{\varphi}(JM) \subset JM + J^2 \otimes V$, and so $\tilde{\varphi}$ induces a T_0 -linear map $\varphi: M/JM \rightarrow JN/J^2N$. Since $T_0 \rightarrow T/J$ is surjective, it follows that φ is T/J -linear as well. It is clear that φ is the unique map satisfying the stated condition. $\square$

PROPOSITION 6.3. *We have a natural isomorphism*

$$\iota: (\mathcal{Q}_1^* \otimes \mathcal{R}_0) \oplus (\mathcal{Q}_0^* \otimes \mathcal{R}_1) \rightarrow \mathcal{J}/\mathcal{J}^2$$

of coherent $\mathcal{O}_{X_{\text{ord}}}$ -modules.

Proof. Since $\mathcal{R}/\mathcal{J}\mathcal{R} \cong \mathcal{R}_0 \oplus \mathcal{R}_1$ and $\mathcal{Q}/\mathcal{J}\mathcal{Q} \cong \mathcal{Q}_0 \oplus \mathcal{Q}_1$, Lemma 6.2 gives a canonical map

$$\mathcal{R}_0 \oplus \mathcal{R}_1 \rightarrow (\mathcal{Q}_0 \oplus \mathcal{Q}_1) \otimes_{\mathcal{O}_X/\mathcal{J}} \mathcal{J}/\mathcal{J}^2,$$

where $\mathcal{R}_0$ maps to $\mathcal{Q}_1$ and $\mathcal{R}_1$ maps to $\mathcal{Q}_0$. We can thus convert it into a map

$$\Phi: (\mathcal{Q}_1^* \otimes \mathcal{R}_0) \oplus (\mathcal{Q}_0^* \otimes \mathcal{R}_1) \rightarrow \mathcal{J}/\mathcal{J}^2.$$

We claim that Φ is an isomorphism.

First, we show that Φ is injective. Note that Φ is $\mathbf{G}_{\text{ord}}$ -equivariant and that $\mathcal{Q}_1^* \otimes \mathcal{R}_0$ and $\mathcal{Q}_0^* \otimes \mathcal{R}_1$ are both irreducible homogeneous bundles which are not isomorphic to each other. So it suffices to show that Φ is not identically 0 on either summand. Using the universal property of $\mathbf{Gr}_{r|s}(V)$, it suffices to give a single superalgebra T together with a rank $r|s$ summand M of $T \otimes V$ such that the pullback of Φ to each summand is non-zero.

This can be done with $T = \mathbf{C}[\epsilon]/(\epsilon^2)$, where ϵ has degree 1. Pick bases $e_1, \dots, e_n$ for V_0 and $f_1, \dots, f_m$ for V_1 . Then we let M be the free T -submodule of $T \otimes V$ with basis elements $v_i = 1 \otimes e_i + \epsilon \otimes f_1$ for $i = 1, \dots, r$ and $w_j = \epsilon \otimes e_1 + 1 \otimes f_j$ for $j = 1, \dots, s$. Then $\varphi(v_i) = \epsilon f_1$ and $\varphi(w_j) = \epsilon e_1$, so that both components are indeed non-zero.

Surjectivity now follows since both sides are equivariant vector bundles of the same dimension. (Note that the rank of $\mathcal{J}/\mathcal{J}^2$ is the odd part of the dimension of X .) $\square$

From the above, we see that we are in the setting of Theorem 2.4. Indeed, X is a smooth supervariety and X_{ord} is projective, and we have a short exact sequence

$$0 \rightarrow \mathcal{J}/\mathcal{J}^2 \rightarrow \epsilon \rightarrow \eta \rightarrow 0,$$

where $\epsilon = W^* \otimes \mathcal{O}_{X_{\text{ord}}}$, with W as in §5. We recall that we are assuming $r \geq s$. From this sequence, one easily sees that $\text{Spec}(\text{Sym}(\eta))$ is identified with the variety Y . Thus the varieties Z and $\tilde{Z}$ appearing in §2.2 and Theorem 2.4 match those studied in §5 in our setting. We thus find the following.

PROPOSITION 6.4. *There exist a natural isomorphism*

$$H^q(X, \text{gr}^{p+q}(\mathcal{O}_X)) = \text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_{p+q}$$

and spectral sequence

$$E_1^{p,q} = \text{Tor}_{-q}^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_p \implies H^{p+q}(X, \mathcal{O}_X).$$

Proof. This follows from Theorem 2.4. The key hypothesis, that the higher cohomology of $\mathcal{O}_Y$ vanishes, is provided by Theorem 5.1(d). $\square$

Recall the super Euler characteristic χ defined in §2.3.

PROPOSITION 6.5. Assume $r \geq s$. We have

$$\chi(\mathbf{Gr}_{r|s}(\mathbf{C}^{n|m})) = \begin{cases} \binom{m}{s} & \text{if } n - m \geq r - s, \\ \binom{n}{s} & \text{if } r = s \text{ and } m > n, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. This follows immediately from Corollary 2.5 and Theorem 5.1. $\square$

6.3 The Lascoux resolution

To make use of Proposition 6.4, we need to know something about the minimal free resolution of $\mathcal{O}_{\tilde{Z}}$: specifically, we need to know its terms, that is, the Tor groups (the differentials will not concern us). Since $\mathcal{O}_{\tilde{Z}}$ is a free $\mathcal{O}_Z$ -module (Theorem 5.1(b)), its Tor groups are direct sums of the Tor groups of $\mathcal{O}_Z$. Lascoux determined the Tor groups of $\mathcal{O}_Z$, and we now review his results.

Let a and b be non-negative integers. Given (integer) partitions α and β with $\ell(\alpha) \leq b$ and $\beta_1 \leq b$, define the partitions

$$\begin{aligned} P_{a,b}(\alpha, \beta) &= (b + \alpha_1, \dots, b + \alpha_b, b^a, \beta_1, \dots, \beta_{\ell(\beta)}), \\ Q_{a,b}(\alpha, \beta) &= (b + \beta_1^\dagger, \dots, b + \beta_b^\dagger, b^a, \alpha_1^\dagger, \dots, \alpha_{\ell(\alpha)}^\dagger), \end{aligned}$$

which we visualize in terms of Young diagrams as follows:

$$P_{a,b}(\alpha, \beta) = \begin{array}{|c|c|} \hline b \times b & \alpha \\ \hline a \times b & \\ \hline \beta & \\ \hline \end{array} \quad Q_{a,b}(\alpha, \beta) = \begin{array}{|c|c|} \hline b \times b & \beta^\dagger \\ \hline a \times b & \\ \hline \alpha^\dagger & \\ \hline \end{array}$$

For a partition λ , we let $\mathbf{S}_\lambda$ denote the corresponding Schur functor. With this notation, we can state Lascoux's result.

THEOREM 6.6 (Lascoux). Put $a = m - \delta$. If $q = ab$ for some non-negative integer b , then

$$\mathrm{Tor}_p^S(\mathcal{O}_Z, \mathbf{C})_{p+q} = \bigoplus_{\substack{\alpha, \beta \\ \ell(\alpha) \leq b, \beta_1 \leq b \\ p = b^2 + |\alpha| + |\beta|}} \mathbf{S}_{P_{a,b}(\alpha, \beta)}(V_0) \otimes \mathbf{S}_{Q_{a,b}(\alpha, \beta)}(V_1^*)$$

as representations of $\mathbf{G}_{\mathrm{ord}}$. If q is not divisible by b , then $\mathrm{Tor}_p^{\mathcal{O}_W}(\mathcal{O}_Z, \mathbf{C})_{p+q} = 0$.

Proof. See [Wey03, Proposition 6.1.3], which contains the description of $\mathrm{Tor}_p^{\mathcal{O}_{W_0}}(\mathcal{O}_{Z_0}, \mathbf{C})$. Note that since $Z = Z_0 \times W_1$, the minimal free resolution for $\mathcal{O}_Z$ over $\mathcal{O}_W$ is obtained from the minimal free resolution for $\mathcal{O}_{Z_0}$ over $\mathcal{O}_{W_0}$ by tensoring with $\mathcal{O}_{W_1}$. In particular, the Tor groups agree, and this conversion preserves the grading. $\square$

COROLLARY 6.7. The $\mathbf{G}_{\mathrm{ord}}$ -representation $\bigoplus_{p \geq 0} \mathrm{Tor}_p^S(\mathcal{O}_Z, \mathbf{C})$ is multiplicity-free.

6.4 Proof of Theorem 6.1

Put

$$\tilde{L}_k = \bigoplus_{p \geq 0} \mathrm{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_{p+k}, \quad L_k = \bigoplus_{p \geq 0} \mathrm{Tor}_p^S(\mathcal{O}_Z, \mathbf{C})_{p+k}.$$

These vector spaces carry algebraic representations of $\mathbf{G}_{\text{ord}}$. In particular, they are semi-simple as $\mathbf{G}_{\text{ord}}$ -representations, as are all algebraic representations.

PROPOSITION 6.8. *The graded vector space $\text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})$ is naturally a graded A -module, and the induced map*

$$A \otimes_{\mathbf{C}} \text{Tor}_p^S(\mathcal{O}_Z, \mathbf{C}) \rightarrow \text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C}) \quad (6.8a)$$

is an isomorphism of graded A -modules.

Proof. The space $\text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})$ is naturally a module over $\text{Tor}_0^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C}) = \mathcal{O}_{\tilde{Z}} \otimes_{\mathcal{O}_Z} \mathbf{C}$, which we have seen (Theorem 5.1(c)) is isomorphic to A . Let $A' \subset \mathcal{O}_{\tilde{Z}}$ be a homogeneous $\mathbf{C}$ -subspace such that the map $A' \rightarrow \mathcal{O}_{\tilde{Z}} \otimes_{\mathcal{O}_Z} \mathbf{C} = A$ is an isomorphism. Then A' is a minimal generating space for $\mathcal{O}_{\tilde{Z}}$ as an $\mathcal{O}_Z$ -module. Since $\mathcal{O}_{\tilde{Z}}$ is free as an $\mathcal{O}_Z$ -module (Theorem 5.1(b)), it follows that the natural map

$$A' \otimes_{\mathbf{C}} \mathcal{O}_Z \rightarrow \mathcal{O}_{\tilde{Z}}$$

is an isomorphism of $\mathcal{O}_Z$ -modules. We thus see that the induced map

$$A' \otimes_{\mathbf{C}} \text{Tor}_p^S(\mathcal{O}_Z, \mathbf{C}) \rightarrow \text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})$$

is an isomorphism. This map is isomorphic (in the obvious manner) to (6.8a), and so (6.8a) is an isomorphism of vector spaces. The map (6.8a) is a homomorphism of graded A -modules simply by its definition. $\square$

PROPOSITION 6.9. *The $\mathbf{G}_{\text{ord}}$ -representations $\tilde{L}_k$ and $\tilde{L}_{k+1}$ have no simple factors in common.*

Proof. Proposition 6.8 shows that $\tilde{L}_k = \bigoplus_{i \geq 0} A_{k-i} \otimes_{\mathbf{C}} L_i$. Since A is concentrated in even degrees, we see that $\tilde{L}_k$ is a sum of L_i with i of the same parity as k . Since $\bigoplus_{i \geq 0} L_i$ is multiplicity-free as a $\mathbf{G}_{\text{ord}}$ -representation (Corollary 6.7), the claim follows. $\square$

PROPOSITION 6.10. *The spectral sequence in Proposition 6.4 degenerates at the first page.*

Proof. Let E be the spectral sequence from Proposition 6.4, and put $E_r^k = \bigoplus_{p+q=k} E_r^{p,q}$, so that the differential is a map $E_r^k \rightarrow E_r^{k+1}$. We have $E_1^k = \tilde{L}_k$. Since E_r^k is a subquotient of E_1^k , we see that E_r^k is a semi-simple $\mathbf{G}_{\text{ord}}$ -representation, and E_r^k and E_r^{k+1} have no irreducible factors in common. It follows that the differential $E_r^k \rightarrow E_r^{k+1}$ must vanish, as it is a map of $\mathbf{G}_{\text{ord}}$ -representations. This completes the proof. $\square$

COROLLARY 6.11. *We have canonical $\mathbf{G}_{\text{ord}}$ -equivariant isomorphisms*

$$H^i(X, \mathcal{O}_X) = \text{gr}(H^i(X, \mathcal{O}_X)) = \tilde{L}_i.$$

Proof. Proposition 6.10 gives a canonical isomorphism $\text{gr}(H^i(X, \mathcal{O}_X)) = \tilde{L}_i$. It follows that $\text{gr}(H^i(X, \mathcal{O}_X))$ is multiplicity-free as a representation of $\mathbf{G}_{\text{ord}}$, and so the same is true of $H^i(X, \mathcal{O}_X)$. Thus the filtration on $H^i(X, \mathcal{O}_X)$ canonically splits, which yields a canonical isomorphism $H^i(X, \mathcal{O}_X) = \text{gr}(H^i(X, \mathcal{O}_X))$. $\square$

PROPOSITION 6.12. *We have $H^i(X, \mathcal{O}_X)^{\mathbf{G}} = H^i(X, \mathcal{O}_X)^{\mathbf{G}_{\text{ord}}}$.*

Proof. First assume $m > \delta$. Let $E = H^i(X, \mathcal{O}_X)^{\mathbf{G}_{\text{ord}}}$. The action of the upper-triangular nilpotent piece of the Lie algebra of $\mathbf{G}$ on W is a $\mathbf{G}_{\text{ord}}$ -equivariant map

$$V_1^* \otimes V_0 \otimes E \rightarrow H^i(X, \mathcal{O}_X).$$

First note that $H^i(X, \mathcal{O}_X)$ does not contain a $\mathbf{G}_{\text{ord}}$ -subrepresentation isomorphic to $V_1^* \otimes V_0$: in the notation of Theorem 6.6, this is only possible if $P_{a,b}(\alpha, \beta) = Q_{a,b}(\alpha, \beta) = (1)$, which is

only possible if $a = m - \delta = 0$. It follows that this map must be 0. Similarly, the corresponding map for the lower-triangular piece is 0. Thus E is annihilated by the Lie algebra of $\mathbf{G}$. It follows that $\mathbf{G}$ acts trivially on E .

Finally, if $m = \delta$, this means that $r = n$ and $s = 0$. In this case, X is a point, and its structure sheaf is the exterior algebra on $V_1^* \otimes V_0$. The $\mathbf{G}_{\text{ord}}$ -invariant space is spanned by the unit element, and since $\mathbf{G}$ acts via algebra automorphisms, the unit must also be invariant under $\mathbf{G}$. $\square$

Let U be a graded vector space. We define the *trivial filtration* on U by $\text{Fil}^i(U) = \bigoplus_{j \geq i} U_j$. With respect to this filtration, we have a natural isomorphism $U = \text{gr}(U)$.

PROPOSITION 6.13. *We have a natural isomorphism $H^*(X, \mathcal{O}_X)^{\mathbf{G}} = A$ of graded algebras. Moreover, the filtration on $H^*(X, \mathcal{O}_X)$ induces the trivial filtration on A .*

Proof. Let $B = H^*(X, \mathcal{O}_X)^{\mathbf{G}}$. By Proposition 6.12, we have $B = H^*(X, \mathcal{O}_X)^{\mathbf{G}_{\text{ord}}}$. We have

$$\begin{aligned} \text{gr}^p(B_q) &= \text{gr}^p(H^q(X, \mathcal{O}_X))^{\mathbf{G}_{\text{ord}}} = H^q(X, \text{gr}^p(\mathcal{O}_X))^{\mathbf{G}_{\text{ord}}} \\ &= \text{Tor}_{p-q}^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_p^{\mathbf{G}_{\text{ord}}} = \bigoplus_{i+j=p} A_i \otimes \text{Tor}_{p-q}^S(\mathcal{O}_Z, \mathbf{C})_j^{\mathbf{G}_{\text{ord}}}. \end{aligned}$$

In the first step, we used that the $\mathbf{G}_{\text{ord}}$ -action is semi-simple; in the second, that the spectral sequence degenerates (Proposition 6.10); in the third, Proposition 6.4; and in the fourth, Proposition 6.8. Now, it is easy to see directly that $\text{Tor}_k^S(\mathcal{O}_Z, \mathbf{C})^{\mathbf{G}_{\text{ord}}}$ vanishes for $k \neq 0$ and that $\text{Tor}_0^S(\mathcal{O}_Z, \mathbf{C})^{\mathbf{G}_{\text{ord}}}$ is 1-dimensional and concentrated in degree 0; this can also be read off of Theorem 6.6. We thus find that

$$\text{gr}^p(B_q) = \begin{cases} A_p & \text{if } p = q, \\ 0 & \text{otherwise.} \end{cases}$$

This shows that $\text{gr}(B) = A$, and the isomorphism is one of rings by Remark 2.3. It also shows that the filtration on B is trivial, and so B is isomorphic to $\text{gr}(B)$ as a ring. $\square$

We regard $H^*(X, \mathcal{O}_X)$ as a graded A -module via Proposition 6.13. We have isomorphisms of $\mathbf{G}_{\text{ord}}$ -representations

$$H^i(X, \mathcal{O}_X) \cong \text{gr}(H^i(X, \mathcal{O}_X)) = \tilde{L}_i = \bigoplus_{j \geq 0} A_{i-j} \otimes L_j.$$

It follows that $H^i(X, \mathcal{O}_X)$ contains a unique $\mathbf{G}_{\text{ord}}$ -subrepresentation isomorphic to L_i . (Corollary 6.7 is important here.) Call this subspace E^i , and let $E = \bigoplus_{i \geq 0} E^i$.

PROPOSITION 6.14. *The natural map $A \otimes E \rightarrow H^*(X, \mathcal{O}_X)$ is an isomorphism.*

Proof. We have

$$\text{gr}(H^*(X, \mathcal{O}_X)) = \bigoplus_{p,q \geq 0} \text{Tor}_p^S(\mathcal{O}_{\tilde{Z}}, \mathbf{C})_q = A \otimes \bigoplus_{p,q \geq 0} \text{Tor}_p(\mathcal{O}_Z, \mathbf{C})_q = A \otimes \bigoplus_{i \geq 0} L_i.$$

We note that the L_i in the final direct sum belongs to $\text{gr}(H^i(X, \mathcal{O}_X))$. We thus see that $\text{gr}(H^*(X, \mathcal{O}_X))$ is a free A -module (this relies on Remark 2.3). Moreover, if $\bar{E}^i \subset \text{gr}(H^i(X, \mathcal{O}_X))$ is the associated graded of E^i , then $\bar{E}^i$ corresponds to the copy of L_i in the final direct sum above. Letting $\bar{E} = \bigoplus_{i \geq 0} \bar{E}^i$, we thus see that the natural map $A \otimes \bar{E} \rightarrow \text{gr}(H^*(X, \mathcal{O}_X))$ is an isomorphism. The result follows from this. $\square$

PROPOSITION 6.15. *Suppose that M and M' are representations of $\mathbf{G}$ such that $M \cong L_i$ and $M' \cong L_j$ as $\mathbf{G}_{\text{ord}}$ -representations, with $i \neq j$. Then $\text{Ext}_{\mathbf{G}}^1(M, M') = 0$.*

Proof. Consider an extension

$$0 \rightarrow M' \rightarrow N \rightarrow M \rightarrow 0.$$

Since $i \neq j$, there is a unique $\mathbf{G}_{\text{ord}}$ -splitting of this sequence. We thus regard M as a $\mathbf{G}_{\text{ord}}$ -subrepresentation of N . Consider the map $V_1^* \otimes V_0 \otimes M \rightarrow N$ giving the action of the upper-triangular nilpotent piece of the Lie algebra of $\mathbf{G}$ on M . Since this map is $\mathbf{G}_{\text{ord}}$ -equivariant, it follows from Theorem 6.6 and Pieri's rule [FH04, Equation (6.8)] that this map is zero. The analog holds for the lower-triangular piece. Thus M is $\mathbf{G}$ -subrepresentation of N , which completes the proof. $\square$

PROPOSITION 6.16. *The subspace E is a $\mathbf{G}$ -subrepresentation of $H^*(X, \mathcal{O}_X)$.*

Proof. Put $H^i = H^i(X, \mathcal{O}_X)$. We prove that E^i is a $\mathbf{G}$ -subrepresentation of H^i by induction on i . Thus suppose that i is given and E^j is a $\mathbf{G}$ -subrepresentation of H^j for all $j < i$. Let $X = \sum_{j=0}^{i-1} A_{i-j} E^j$, which is a $\mathbf{G}$ -subrepresentation of H^i by the inductive hypothesis, and let $Y = H^i/X$. We thus have a short exact sequence

$$0 \rightarrow X \rightarrow H^i \rightarrow Y \rightarrow 0$$

of $\mathbf{G}$ -representations. As the map $A \otimes E \rightarrow H^*$ is an isomorphism, we see that $X \cong \bigoplus_{j=0}^{i-1} A_{i-j} \otimes E^j$ and that E^i is a complementary subspace to X in H^i . In particular, the map $E^i \rightarrow Y$ is an isomorphism of $\mathbf{G}_{\text{ord}}$ -representations. It now follows from Proposition 6.15 that $\text{Ext}_{\mathbf{G}}^1(Y, X) = 0$. Indeed, we have $E^j \cong L_j$ and $Y \cong L_i$ as $\mathbf{G}_{\text{ord}}$ -representations and $i \neq j$. We thus see that there is a $\mathbf{G}$ -splitting $\sigma: Y \rightarrow H^i$. Since E^i is the unique $\mathbf{G}_{\text{ord}}$ -subrepresentation of H^i isomorphic to L_i , we see that σ is unique and $\sigma(Y) = E^i$. Thus E^i is a $\mathbf{G}$ -subrepresentation, as required. $\square$

6.5 The $\mathbf{G}$ -action on the Lascoux resolution

Our original motivation for this work was to give a geometric explanation for the result of [PW85] (and [Sam14]) that the linear strands of the Tor groups of the determinantal variety Z_0 carry an action of $\mathbf{G}$. Of course, it is equivalent to do this for Z since Z and Z_0 have the same Tor groups. This follows from Proposition 6.16 as $E^i = L_i$ is a $\mathbf{G}$ -subrepresentation of $H^i(X, \mathcal{O}_X)$. If one is simply interested in constructing the $\mathbf{G}$ -action on L_i , it suffices to consider the case where $s = 0$; here we have

$$L_i = H^i(X, \mathcal{O}_X),$$

and so the obvious $\mathbf{G}$ -action on the right induces one on the left.

REMARK 6.17. The action of $\mathbf{G}$ of L_i induces an action of its Lie superalgebra on L_i . With respect to the decomposition $\mathfrak{gl}(n|m) = \text{Hom}(\mathbf{C}^n, \mathbf{C}^m) \oplus (\mathfrak{gl}(n) \times \mathfrak{gl}(m)) \oplus \text{Hom}(\mathbf{C}^m, \mathbf{C}^n)$, the middle piece corresponds to the obvious action of $\mathbf{GL}(n) \times \mathbf{GL}(m)$ on determinantal varieties, while one of the other pieces (depending on conventions) corresponds to the action of the Tor algebra on linear strands by Remark 2.3. With that structure in place, the L_i are highest weight representations of $\mathbf{G}$, and if $m > \delta$, the results of [PW85, Sam14] show that they are irreducible (and hence there is a unique way to extend the action of two out of the three pieces to the third). If $m = \delta$, then L_i is only non-zero for $i = 0$, in which case it is a Koszul complex, which we can understand as a Kac module, that is, induced from the trivial representation of $\text{Hom}(\mathbf{C}^n, \mathbf{C}^m) \oplus (\mathfrak{gl}(n) \times \mathfrak{gl}(m))$ (again depending on conventions). It is indecomposable, but not irreducible in general.

REFERENCES

- AW97 K. Akin and J. Weyman, *Minimal free resolutions of determinantal ideals and irreducible representations of the Lie superalgebra $\mathfrak{gl}(m|n)$* , J. Algebra **197** (1997), no. 2, 559–583; doi:[10.1006/jabr.1997.7101](https://doi.org/10.1006/jabr.1997.7101).
- AW00 ———, *The irreducible tensor representations of $\mathfrak{gl}(m|1)$ and their generic homology*, J. Algebra **230** (2000), no. 1, 5–23; doi:[10.1006/jabr.1999.7986](https://doi.org/10.1006/jabr.1999.7986).
- AW07 ———, *Primary ideals associated to the linear strands of Lascoux’s resolution and syzygies of the corresponding irreducible representations of the Lie superalgebra $\mathfrak{gl}(m|n)$* , J. Algebra **310** (2007), no. 2, 461–490; doi:[10.1016/j.jalgebra.2003.11.015](https://doi.org/10.1016/j.jalgebra.2003.11.015).
- Cou16 K. Coulembier, *Bott–Borel–Weil theory and Bernstein–Gel’fand–Gel’fand reciprocity for Lie superalgebras*, Transform. Groups **21** (2016), no. 3, 681–723; doi:[10.1007/s00031-016-9377-6](https://doi.org/10.1007/s00031-016-9377-6).
- DW17 H. Derksen and J. Weyman, *An Introduction to Quiver Representations*, Grad. Stud. Math., vol. 184 (Amer. Math. Soc., Providence, RI), 2017; doi:[10.1090/gsm/184](https://doi.org/10.1090/gsm/184).
- EL05 T. Ekedahl and D. Laksov, *Splitting algebras, symmetric functions and Galois theory*, J. Algebra Appl. **4** (2005), no. 1, 59–75; doi:[10.1142/S0219498805001034](https://doi.org/10.1142/S0219498805001034).
- FH04 W. Fulton and J. Harris, *Representation Theory*, Grad. Stud. Math., vol. 129 (Springer-Verlag, New York, 2004); doi:[10.1007/978-1-4612-0979-9](https://doi.org/10.1007/978-1-4612-0979-9).
- GSS12 D. R. Grayson, A. Seceleanu, and M. E. Stillman, *Computations in intersection rings of flag bundles*, 2012, [arXiv:1205.4190](https://arxiv.org/abs/1205.4190).
- GS10 C. Gruson and V. Serganova, *Cohomology of generalized supergrassmannians and character formulae for basic classical Lie superalgebras*, Proc. Lond. Math. Soc. (3) **101** (2010), no. 3, 852–892; doi:[10.1112/plms/pdq014](https://doi.org/10.1112/plms/pdq014).
- Har94 R. Hartshorne, *Generalized divisors on Gorenstein schemes*, Proceedings Conference on Algebraic Geometry and Ring Theory in honor of Michael Artin, Part III (Antwerp, 1992), K-Theory **8** (1994), no. 3, 287–339; doi:[10.1007/BF00960866](https://doi.org/10.1007/BF00960866).
- KM98 J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., vol. 134 (Cambridge Univ. Press, Cambridge, 1998); doi:[10.1017/CB09780511662560](https://doi.org/10.1017/CB09780511662560).
- Lak10 D. Laksov, *Splitting algebras and Gysin homomorphisms*, J. Commut. Algebra **2** (2010), no. 3, 401–425; doi:[10.1216/JCA-2010-2-3-401](https://doi.org/10.1216/JCA-2010-2-3-401).
- Las78 A. Lascoux, *Syzygies des variétés déterminantales*, Adv. Math. **30** (1978), no. 3, 202–237; doi:[10.1016/0001-8708\(78\)90037-3](https://doi.org/10.1016/0001-8708(78)90037-3).
- Man97 Yu. I. Manin, *Gauge Field Theory and Complex Geometry*, 2nd ed., Grundlehren math. Wiss., vol. 289 (Springer-Verlag, Berlin, 1997); doi:[10.1007/978-3-662-07386-5](https://doi.org/10.1007/978-3-662-07386-5).
- MZ11 A. Masuoka and A. N. Zubkov, *Quotient sheaves of algebraic supergroups are superschemes*, J. Algebra **348** (2011), 135–170; doi:[10.1016/j.jalgebra.2011.08.038](https://doi.org/10.1016/j.jalgebra.2011.08.038).
- MS11 I. M. Musson and V. Serganova, *Combinatorics of character formulas for the Lie superalgebra $\mathfrak{gl}(m, n)$* , Transform. Groups **16** (2011), no. 2, 555–578; doi:[10.1007/s00031-011-9147-4](https://doi.org/10.1007/s00031-011-9147-4).
- NSS22 R. Nagpal, S. V. Sam, and A. Snowden, *On the geometry and representation theory of isomeric matrices*, Algebra Number Theory **16** (2022), no. 6, 1501–1529; doi:[10.2140/ant.2022.16.1501](https://doi.org/10.2140/ant.2022.16.1501).
- Pen90 I. B. Penkov, *Borel–Weil–Bott theory for classical Lie superalgebras*, J. Soviet Math. **51** (1990), 2108–2140; doi:[10.1007/BF01098186](https://doi.org/10.1007/BF01098186).
- PS97 I. Penkov and V. Serganova, *Characters of irreducible G -modules and cohomology of G/P for the Lie supergroup $G = Q(N)$* , J. Math. Sci. **84** (1997), no. 5, 1382–1412; doi:[10.1007/BF02399196](https://doi.org/10.1007/BF02399196).
- PW85 P. Pragacz and J. Weyman, *Complexes associated with trace and evaluation. Another approach to Lascoux’s resolution*, Adv. Math. **57** (1985), no. 2, 163–207; doi:[10.1016/0001-8708\(85\)90052-0](https://doi.org/10.1016/0001-8708(85)90052-0).
- RW19 C. Raicu and J. Weyman, *Syzygies of determinantal thickenings and representations of the general linear Lie superalgebra*, Acta Math. Vietnam. **44** (2019), no. 1, 269–284; doi:[10.1007/s40306-018-0282-z](https://doi.org/10.1007/s40306-018-0282-z).

- Sam14 S. V Sam, *Derived supersymmetries of determinantal varieties*, J. Commut. Algebra **6** (2014), no. 2, 261–286; doi:[10.1216/JCA-2014-6-2-261](https://doi.org/10.1216/JCA-2014-6-2-261).
- Sam16 ———, *Orthosymplectic Lie superalgebras, Koszul duality, and a complete intersection analogue of the Eagon–Northcott complex*, J. Eur. Math. Soc. **18** (2016), no. 12, 2691–2732; doi:[10.4171/JEMS/651](https://doi.org/10.4171/JEMS/651).
- Ser96 V. Serganova, *Kazhdan–Lusztig polynomials and character formula for the Lie superalgebra $\mathfrak{gl}(m|n)$* , Selecta Math. (N.S.) **2** (1996), no. 4, 607–651; doi:[10.1007/PL00001385](https://doi.org/10.1007/PL00001385).
- Sta21 The Stacks project authors, *The Stacks Project*, 2021, <https://stacks.math.columbia.edu>.
- Wey03 J. Weyman, *Cohomology of Vector Bundles and Syzygies*, Cambridge Tracts in Math., vol. 149 (Cambridge Univ. Press, Cambridge, 2003); doi:[10.1017/CB09780511546556](https://doi.org/10.1017/CB09780511546556).

Steven V Sam ssam@ucsd.edu

Department of Mathematics, University of California, San Diego, CA, USA

Andrew Snowden asnowden@umich.edu

Department of Mathematics, University of Michigan, Ann Arbor, MI, USA